

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 June 2006 (15.06.2006)

PCT

(10) International Publication Number
WO 2006/062465 A1

(51) International Patent Classification:
C07D 487/04 (2006.01) **A61P 9/10** (2006.01)
A61K 31/519 (2006.01)

(74) Agent: **ASTRAZENECA**; Global Intellectual Property,
S-151 85 Södertälje (SE).

(21) International Application Number:
PCT/SE2005/001835

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
5 December 2005 (05.12.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0402972-4 6 December 2004 (06.12.2004) SE
0501093-9 13 May 2005 (13.05.2005) SE

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **ASTRAZENECA AB** [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **BÖGEVIG, Anders** [SE/SE]; AstraZeneca R & D Södertälje, S-151 85 Södertälje (SE). **LO-ALFREDSSON, Yvonne** [SE/SE]; AstraZeneca R & D Södertälje, S-151 85 Södertälje (SE). **PIVONKA, Donald** [US/US]; AstraZeneca Wilmington, 1800 Concord Pike, P.O. Box 15437, Wilmington, DE 19850-5437 (US). **TIDÉN, Anna-Karin** [SE/SE]; AstraZeneca R & D Södertälje, S-151 85 Södertälje (SE).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL PYRROLO [3, 2-D] PYRIMIDIN-4-ONE DERIVATIVES AND THEIR USE IN THERAPY



(I)

(57) Abstract: There are disclosed novel compounds of formula (I) wherein R¹, R¹², L, X and Y are as defined in the specification, and pharmaceutically acceptable salts thereof; together with processes for their preparation, compositions containing them and their use in therapy. The compounds are inhibitors of the enzyme MPO and are thereby particularly useful in the treatment or prophylaxis of neuroinflammatory disorders, cardiovascular disorders and respiratory disorders.

WO 2006/062465 A1

Novel pyrrolo[3,2-d]pyrimidin-4-one derivatives and their use in therapy.

Field of the invention

The present invention relates to novel pyrrolo[3,2-d]pyrimidin-4-one derivatives, processes
5 for their preparation, compositions containing them and their use in therapy.

Background of the invention

Myeloperoxidase (MPO) is a heme-containing enzyme found predominantly in polymorphonuclear leukocytes (PMNs). MPO is one member of a diverse protein family of
10 mammalian peroxidases that also includes eosinophil peroxidase, thyroid peroxidase, salivary peroxidase, lactoperoxidase, prostaglandin H synthase, and others. The mature enzyme is a dimer of identical halves. Each half molecule contains a covalently bound heme that exhibits unusual spectral properties responsible for the characteristic green colour of MPO. Cleavage of the disulphide bridge linking the two halves of MPO yields
15 the hemi-enzyme that exhibits spectral and catalytic properties indistinguishable from those of the intact enzyme. The enzyme uses hydrogen peroxide to oxidize chloride to hypochlorous acid. Other halides and pseudohalides (like thiocyanate) are also physiological substrates to MPO.

20 PMNs are of particular importance for combating infections. These cells contain MPO, with well documented microbicidal action. PMNs act non-specifically by phagocytosis to engulf microorganisms, incorporate them into vacuoles, termed phagosomes, which fuse with granules containing myeloperoxidase to form phagolysosomes. In phagolysosomes the enzymatic activity of the myeloperoxidase leads to the formation of hypochlorous acid,
25 a potent bactericidal compound. Hypochlorous acid is oxidizing in itself, and reacts most avidly with thiols and thioethers, but also converts amines into chloramines, and chlorinates aromatic amino acids. Macrophages are large phagocytic cells which, like PMNs, are capable of phagocytosing microorganisms. Macrophages can generate hydrogen peroxide and upon activation also produce myeloperoxidase. MPO and hydrogen

peroxide can also be released to the outside of the cells where the reaction with chloride can induce damage to adjacent tissue.

Linkage of myeloperoxidase activity to disease has been implicated in neurological diseases with a neuroinflammatory response including multiple sclerosis, Alzheimer's disease, Parkinson's disease and stroke as well as other inflammatory diseases or conditions like asthma, chronic obstructive pulmonary disease, cystic fibrosis, atherosclerosis, inflammatory bowel disease, renal glomerular damage and rheumatoid arthritis. Lung cancer has also been suggested to be associated with high MPO levels.

Multiple sclerosis (MS)

MPO positive cells are immensely present in the circulation and in tissue undergoing inflammation. More specifically MPO containing macrophages and microglia has been documented in the CNS during disease; multiple sclerosis (Nagra RM, et al. Journal of Neuroimmunology 1997; 78(1-2):97-107), Parkinson's disease (Choi D-K. et al. J. Neurosci. 2005; 25(28):6594-600) and Alzheimer's disease (Green PS. et al. Journal of Neurochemistry. 2004; 90(3):724-33). It is supposed that some aspects of a chronic ongoing inflammation result in an overwhelming destruction where agents from MPO reactions have an important role.

The enzyme is released both extracellularly as well as into phagolysosomes in the neutrophils (Hampton MB, Kettle AJ, Winterbourn CC. Blood 1998; 92(9):3007-17). A prerequisite for the MPO activity is the presence of hydrogen peroxide, generated by NADPH oxidase and a subsequent superoxide dismutation. The oxidized enzyme is capable to use a plethora of different substrates of which chloride is most recognized. From this reaction the strong non-radical oxidant - hypochlorous acid (HOCl) - is formed. HOCl oxidizes sulphur containing amino acids like cysteine and methionine very efficiently (Peskin AV, Winterbourn CC. Free Radical Biology and Medicine 2001; 30(5):572-9). It also forms chloramines with amino groups, both in proteins and other biomolecules (Peskin AV. et al. Free Radical Biology and Medicine 2004; 37(10):1622-30). It

chlorinates phenols (like tyrosine) (Hazen SL. et al. Mass Free Radical Biology and Medicine 1997; 23(6):909-16) and unsaturated bonds in lipids (Albert CJ. et al. J. Biol. Chem. 2001; 276(26):23733-41), oxidizes iron centers (Rosen H, Klebanoff SJ. Journal of Biological Chemistry 1982; 257(22):13731-354) and crosslinks proteins (Fu X, Mueller DM, Heinecke JW. Biochemistry 2002; 41(4):1293-301).

Proteolytic cascades participate both in cell infiltration through the BBB as well as the destruction of BBB, myelin and nerve cells (Cuzner ML, Opdenakker G. Journal of Neuroimmunology 1999; 94(1-2):1-14; Yong VW. et al. Nature Reviews Neuroscience 2001; 2(7):502-11.). Activation of matrix metalloproteinases (MMPs) can be accomplished through the action of upstream proteases in a cascade as well as through oxidation of a disulfide bridge (Fu X. et al. J. Biol. Chem. 2001; 276(44):41279-87; Gu Z. et al. Science 2002; 297(5584):1186-90). This oxidation can be either a nitrosylation or HOCl-mediated oxidation. Both reactions can be a consequence of MPO activity. Several reports have suggested a role for MMP's in general and MMP-9 in particular as influencing cell infiltration as well as tissue damage (BBB breakdown and demyelination), both in MS and EAE (for review see Yong VW. et al, *supra*). The importance of these specific kinds of mechanisms in MS comes from studies where increased activity and presence of proteases have been identified in MS brain tissue and CSF. Supportive data has also been generated by doing EAE studies with mice deficient in some of the proteases implicated to participate in the MS pathology, or by using pharmacological approaches.

The demyelination is supposed to be dependent on the cytotoxic T-cells and toxic products generated by activated phagocytes (Lassmann H. J Neurol Neurosurg Psychiatry 2003; 74(6):695-7). The axonal loss is thus influenced by proteases and reactive oxygen and nitrogen intermediates. When MPO is present it will obviously have the capability of both activating proteases (directly as well as through disinhibition by influencing protease inhibitors) and generating reactive species.

Chronic obstructive pulmonary disease (COPD)

Chronic obstructive pulmonary disease (COPD) is a disease state characterised by airflow limitation that is not fully reversible. The airflow limitation is usually both progressive and associated with an abnormal inflammatory response of the lungs to noxious particles or gases. COPD is a major public health problem. It is the fourth leading cause of chronic morbidity and mortality in the United States¹ and is projected to rank fifth in 2020 as a worldwide burden of disease. In the UK the prevalence of COPD is 1.7% in men and 1.4% in women. COPD spans a range of severity from mild to very severe, with the cost of treatment rising rapidly as the severity increases.

Levels of MPO in sputum and BAL are much greater in COPD patients than normal, non-smoking controls (Keatings V.M., Barnes P.J. *Am J Respir Crit Care Med* 1997;155:449-453; Pesci, A. et al. *Eur Respir J* 1998;12:380-386). MPO levels are further elevated during exacerbations of the disease (Fiorini G. et al. *Biomedicine & Pharmacotherapy* 2000; 54:274-278; Crooks S.W. et al. *European Respiratory Journal*. 15(2):274-80, 2000). The role of MPO is likely to be more important in exacerbations of COPD (Sharon S.D. et al. *Am J Respir Crit Care Med*. 2001;163:349-355).

In addition to the destructive capacity of MPO there is a strong clinical link with vascular disease (Baldus S. et al. *Circulation* 2003;108:1440-5). Dysfunctional MPO polymorphisms are associated with a reduced risk of mortality from coronary artery disease (Nikpoor B. et al. *Am Heart J* 2001; 142:336), and patients with high serum levels of MPO have increased risk of acute coronary syndrome. The effects of MPO on vascular disease may extend to COPD, since there is strong evidence that the pulmonary vasculature is one of the earliest sites of involvement in the smokers' lung. Striking changes in the intima of the pulmonary arteries have been described which show a dose relationship with smoking (Hale K.A., Niewoehner D.E., Cosio M.G. *Am Rev Resp Dis* 1980;122:273-8). The physiological function of MPO is associated with innate host defence. This role, however, is not critical as most cases of MPO deficient patients have relatively benign symptoms (Parry M.F. et al. *Ann Int Med*. 1981; 95:293-301, Yang, K.D., Hill, H.R. *Pediatr Infect Dis J*. 2001;20: 889-900). In summary, there is considerable evidence that


elevated MPO levels in COPD may contribute to the disease via several mechanisms. A selective inhibitor of MPO would therefore be expected to alleviate both the acute and chronic inflammatory aspects of COPD and may reduce the development of emphysema.

5 *Atherosclerosis*

An MPO inhibitor should reduce the atherosclerotic burden and/or the vulnerability of existing atherosclerotic lesions and thereby decrease the risk of acute myocardial infarction, unstable angina or stroke. Several lines of data support a role for MPO in atherosclerosis. MPO is expressed in the shoulder regions and necrotic core of human
10 atherosclerotic lesions and active enzyme has been isolated from autopsy specimens of human lesions (Daugherty, A. et al. (1994) J Clin Invest 94(1): 437-44). In eroded and ruptured human lesions, as compared to fatty streaks, an increased number of MPO expressing macrophages have been demonstrated, suggesting a particular role for MPO in acute coronary syndromes (Sugiyama, S. et al. (2001) Am J Pathol 158(3): 879-91).
15 Patients with established coronary artery disease have higher plasma and leukocyte MPO levels than healthy controls (Zhang, R. et al. (2001) Jama 286(17): 2136-42). Moreover, in two large prospective studies plasma levels of MPO predicted the risk of future coronary events or revascularisation (Baldus, S. et al. (2003) Circulation 108(12): 1440-5; Brennan, M. et al. (2003) N Engl J Med 349(17): 1595-604). Total MPO deficiency in humans has a
20 prevalence of 1 in 2000-4000 individuals. These individuals appear principally healthy but a few cases of severe Candida infection have been reported. Interestingly, MPO deficient humans are less affected by cardiovascular disease than controls with normal MPO levels (Kutter, D. et al. (2000) Acta Haematol 104(1)). A polymorphism in the MPO promoter affects expression leading to high and low MPO expressing individuals. In three different
25 studies the high expression genotype has been associated with an increased risk of cardiovascular disease (Nikpoor, B. et al. (2001) Am Heart J 142(2): 336-9; Makela, R., P. J. Karhunen, et al. (2003) Lab Invest 83(7): 919-25; Asselbergs, F. W., et al. (2004) Am J Med 116(6): 429-30). Data accumulated during the last ten years indicate that the proatherogenic actions of MPO include oxidation of lipoproteins, induction of endothelial
30 dysfunction via consuming nitric oxide and destabilisation of atherosclerotic lesions by


activation of proteases (Nicholls, S. J. and S. L. Hazen (2005) *Arterioscler Thromb Vasc Biol* 25(6): 1102-11). Recently, several studies have focused on nitro- and chlorotyrosine modifications of LDL and HDL lipoproteins. Since chlorotyrosine modifications in vivo only can be generated by hypochlorous acid produced by MPO these modifications are regarded as specific markers of MPO activity (Hazen, S. L. and J. W. Heinecke (1997) *J Clin Invest* 99(9): 2075-81). LDL particles exposed to MPO in vitro become aggregated, leading to facilitated uptake via macrophage scavenger receptors and foam cell formation (Hazell, L. J. and R. Stocker (1993) *Biochem J* 290 (Pt 1): 165-72). Chlorotyrosine modification of apoA1, the main apolipoprotein of HDL cholesterol, results in impaired cholesterol acceptor function (Bergt, C., S. et al. (2004) *Proc Natl Acad Sci U S A*; Zheng, L. et al. (2004) *J Clin Invest* 114(4): 529-41). Systematic studies of these mechanisms have shown that MPO binds to and travels with apoA1 in plasma. Moreover, MPO specifically targets those tyrosine residues of apoA1 that physically interact with the macrophage ABCA1 cassette transporter during cholesterol efflux from the macrophage (Bergt, C. et al. (2004) *J Biol Chem* 279(9): 7856-66; Shao, B. et al. (2005) *J Biol Chem* 280(7): 5983-93; Zheng et al. (2005) *J Biol Chem* 280(1): 38-47). Thus, MPO seems to have a dual aggravating role in atherosclerotic lesions, i.e. increasing lipid accumulation via aggregation of LDL particles and decreasing the reverse cholesterol transport via attack on the HDL protein apoA1.

1- β -D-Ribofuranosyl-2-oxopyrrolo[3,2-d]pyrimidine-4(3H,5H)-thione and 1-(2,3,5-tri-*O*-benzoyl-1- β -D-ribofuranosyl)-2-oxopyrrolo[3,2-d]pyrimidine-4(3H,5H)-thione



are disclosed in J. Heterocyclic Chemistry, 1992, 29, 343-354. No pharmacological activity is ascribed to these compounds.

5 5,7-Dimercapto-1,4,6-triazaindene



is disclosed in Chem. Pharm. Bull., 1964, 12, 1030-1042 and in Japanese patent JP 02160235 A2. No pharmacological activity is ascribed to this compound.


10

The present invention discloses novel pyrrolo[3,2-d]pyrimidin-4-one derivatives that surprisingly display useful properties as inhibitors of the enzyme MPO. These compounds may also show selectivity against related enzymes e.g. lactoperoxidase (LPO) and thyroidperoxidase (TPO).

15

Disclosure of the invention

The present invention provides compounds of formula (I)



5 wherein:

at least one of X and Y represents S, and the other represents O or S;

L represents a direct bond or C1 to 7 alkylene, said alkylene optionally incorporating a
heteroatom selected from O, S(O)_n and NR⁶, said alkylene optionally incorporating one or
10 two carbon-carbon double bonds, and said alkylene being optionally substituted by one or
more substituents selected independently from OH, halogen, CN and NR⁴R⁵, C1 to 6 alkyl
and C1 to 6 alkoxy, said alkoxy optionally incorporating a carbonyl adjacent to the
oxygen;

15 n represents an integer 0, 1 or 2;

R¹ represents hydrogen, or

i) a saturated or partially unsaturated 3 to 7 membered ring optionally incorporating one
20 or two heteroatoms selected independently from O, N and S, and optionally incorporating a
carbonyl group, optionally substituted by one or more substituents independently selected
from halogen, SO₂R⁹, SO₂NR⁹R¹⁰, OH, C1 to 7 alkyl, C1 to 7 alkoxy, CN, CONR²R³,
NR²COR³ and COR³, said alkoxy being optionally further substituted by C1 to 6 alkoxy
and said alkoxy optionally incorporating a carbonyl adjacent to the oxygen, and said alkyl
25 being optionally further substituted by hydroxy or C1 to 6 alkoxy and said alkyl or alkoxy

optionally incorporating a carbonyl adjacent to the oxygen or at any position in the alkyl;
or

ii) an aromatic ring system selected from phenyl, biphenyl, naphthyl or a monocyclic or
5 bicyclic heteroaromatic ring structure containing 1 to 3 heteroatoms independently selected
from O, N and S, said aromatic ring system being optionally substituted by one or more
substituents independently selected from halogen, SO_2R^9 , $\text{SO}_2\text{NR}^9\text{R}^{10}$, OH, C1 to 7 alkyl,
C1 to 7 alkoxy, CN, CONR^2R^3 , NR^2COR^3 and COR^3 ; said alkoxy being optionally
further substituted by C1 to 6 alkoxy and said alkoxy optionally incorporating a carbonyl
10 adjacent to the oxygen, and said alkyl being optionally further substituted by hydroxy or
C1 to 6 alkoxy and said alkyl or alkoxy optionally incorporating a carbonyl adjacent to the
oxygen or at any position in the alkyl;

R^{12} represents hydrogen or halogen or a carbon optionally substituted with one to three
15 halogen atoms;

at each occurrence, R^2 , R^3 , R^4 , R^5 , R^6 , R^9 and R^{10} independently represent hydrogen, C1
to 6 alkyl or C1 to 6 alkoxy said alkoxy optionally incorporating a carbonyl adjacent to the
oxygen, said alkyl being optionally further substituted by halogen, C1 to 6 alkoxy, CHO,
20 C2 to 6 alkanoyl, OH, CONR^7R^8 and NR^7COR^8 ;

or the groups NR^2R^3 , NR^4R^5 and NR^9R^{10} each independently represent a 5 to 7
membered saturated azacyclic ring optionally incorporating one additional heteroatom
selected from O, S and NR^{11} , said ring being optionally further substituted by halogen, C1
25 to 6 alkoxy, CHO, C2 to 6 alkanoyl, OH, CONR^7R^8 and NR^7COR^8 ;

at each occurrence R^7 , R^8 and R^{11} independently represent hydrogen or C1 to 6 alkyl, or
the group NR^7R^8 represents a 5- to 7-membered saturated azacyclic ring optionally
incorporating one additional heteroatom selected from O, S and NR^{11} ;

and pharmaceutically acceptable salts thereof;
with the proviso that the compounds 1- β -D-ribofuranosyl-2-oxopyrrolo[3,2-d]pyrimidine-4(3H,5H)-thione, 1-(2,3,5-tri-*O*-benzoyl-1- β -D-ribofuranosyl)-2-oxopyrrolo[3,2-d]pyrimidine-4(3H,5H)-thione and 5,7-dimercapto-1,4,6-triazaindene are disclaimed.

5

The compounds of formula (I) may exist in enantiomeric forms. It is to be understood that all enantiomers, diastereomers, racemates, tautomers and mixtures thereof are included within the scope of the invention.

10 The compounds of formula (I) may exist in tautomeric forms. All such tautomers and mixtures of tautomers are included within the scope of the present invention.

Unless otherwise indicated, the term "C1 to 6 alkyl" referred to herein denotes a straight or branched chain alkyl group having from 1 to 6 carbon atoms. Examples of such groups
15 include, but are not limited to, methyl, ethyl, 1-propyl, n-butyl, iso-butyl, tert-butyl, pentyl and hexyl. The term "C1 to 7 alkyl" is to be interpreted analogously

Unless otherwise indicated, the term "C1 to 7 alkylene" referred to herein denotes a straight or branched chain alkyl group having from 1 to 7 carbon atoms having two free
20 valencies. Examples of such groups include, but are not limited to, methylene, ethylene, propylene, hexamethylene and ethylethylene. The term "C1 to 3 alkylene" is to be interpreted analogously.

Unless otherwise indicated, the term "C1 to 6 alkoxy" referred to herein denotes a straight
25 or branched chain alkoxy group having from 1 to 6 carbon atoms. Examples of such groups include, but are not limited to, methoxy, ethoxy, 1-propoxy, 2-propoxy (iso-propoxy), tert-butoxy and pentoxy. The term "C1 to 7 alkoxy" is to be interpreted analogously.

Unless otherwise indicated, the term "C2 to 6 alkanoyl" referred to herein denotes a straight or branched chain alkyl group having from 1 to 5 carbon atoms with optional position on the alkyl group by a carbonyl group. Examples of such groups include, but are not limited to, acetyl, propionyl and pivaloyl.

5

Unless otherwise indicated, the term "halogen" referred to herein denotes fluoro, chloro, bromo and iodo.

10 Examples of a saturated or partially unsaturated 3- to 7-membered ring optionally incorporating one or two heteroatoms selected independently from O, N and S, and optionally incorporating a carbonyl group includes, but is not limited to, cyclopropane, cyclopentane, cyclohexane, cyclohexene, cyclopentanone, tetrahydrofuran, pyrrolidine, piperidine, tetrahydropyridine, morpholine, piperazine, pyrrolidinone and piperidinone.

15 Examples of a monocyclic or bicyclic heteroaromatic ring structure containing 1 to 3 heteroatoms independently selected from O, N and S includes, but is not limited to, furan, thiophene, pyrrole, oxazole, isoxazole, thiazole, imidazole, pyrazole, triazole, tetrazole, pyridine, pyrazine, pyrimidine, pyridazine, benzofuran, indole, isoindole and benzimidazole.

20

Examples of a 5 to 7 membered saturated azacyclic ring optionally incorporating one additional heteroatom selected from O, S and NR^{11} includes, but is not limited to, pyrrolidine, piperidine, piperazine, morpholine and thiomorpholine.

25 In the definition of L, "C1 to 7 alkylene; said alkylene optionally incorporating a heteroatom selected from O, S(O)_n and NR^6 ; said alkylene optionally incorporating one or two carbon-carbon double bonds" embraces a saturated or unsaturated straight or branched chain arrangement of 1 to 7 carbon atoms having two free valencies and in which any two singly bonded carbon atoms are optionally separated by O, S or NR^6 . The definition thus
30 includes, for example, methylene, ethylene, propylene, hexamethylene, ethylethylene, $-\text{CH}_2=\text{CH}_2-$, $-\text{CH}_2\text{CH}=\text{CH}-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)=\text{CH}_2-$, $-\text{CH}_2=\text{CH}_2-\text{CH}_2\text{O}-$, $-\text{CH}_2\text{O}-$,

$-\text{CH}_2\text{CH}_2\text{O}-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{O}-\text{CH}_2-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{S}-$ and $-\text{CH}_2\text{CH}_2\text{NR}^6-$.

In one embodiment, R^1 represents hydrogen.

5 In another embodiment, X represents S and Y represents O.

In yet another embodiment, Y represents S and X represents O.

In yet another embodiment, L is a direct bond or represents C1 to 7 alkylene, said alkylene
10 optionally incorporating a heteroatom selected from O, $\text{S}(\text{O})_n$ and NR^6 , said alkylene
optionally incorporating one or two carbon-carbon double bonds, and said alkylene being
optionally substituted by one or more substituents selected independently from OH, C1 to
6 alkoxy, halogen, CN and NR^4R^5 .

15 In yet another embodiment, L is a direct bond or represents C1 to 7 alkylene; said alkylene
being optionally substituted by one or more substituents selected independently from OH,
C1 to 6 alkoxy, halogen, CN and NR^4R^5 .

In yet another embodiment, L is a direct bond or represents C1 to 7 alkylene; said alkylene
20 being optionally substituted by one or more C1 to 6 alkoxy.

In yet another embodiment, L is a direct bond or represents C1 to 3 alkylene; said alkylene
being optionally substituted by one or more substituents selected independently from OH,
C1 to 6 alkoxy, halogen, CN and NR^4R^5 .

25

In yet another embodiment, L represents C1 to 3 alkylene; said alkylene being optionally
substituted by one or more C1 to 6 alkoxy.

In yet another embodiment, L is a direct bond or represents optionally substituted
30 methylene ($-\text{CH}_2-$).

In yet another embodiment, L is a direct bond or represents optionally substituted ethylene (-CH₂CH₂-).

- 5 In yet another embodiment, R¹ represents a saturated or partially unsaturated 3 to 7 membered ring optionally incorporating one or two heteroatoms selected independently from O, N and S, and optionally incorporating a carbonyl group, said ring being optionally substituted by one or more substituents independently selected from halogen, SO₂R⁹,
SO₂NR⁹R¹⁰, OH, C1 to 6 alkyl, C1 to 6 alkoxy, CN, CONR²R³, NR²COR³ and COR³,
10 said alkoxy being optionally further substituted by C1 to 6 alkoxy; and said alkyl being optionally further substituted by hydroxy or C1 to 6 alkoxy.

- In yet another embodiment, R¹ represents a saturated or partially unsaturated 3 to 7 membered ring optionally incorporating one or two heteroatoms selected independently
15 from O, N and S, and optionally incorporating a carbonyl group; said ring being optionally substituted by one or more substituents independently selected from halogen, C1 to 6 alkyl and C1 to 6 alkoxy, said alkoxy being optionally further substituted by C1 to 6 alkoxy.

- In yet another embodiment, R¹ represents an aromatic ring system selected from phenyl,
20 biphenyl, naphthyl or a monocyclic or bicyclic heteroaromatic ring structure containing 1 to 3 heteroatoms independently selected from O, N and S, said aromatic ring being optionally substituted by one or more substituents independently selected from halogen, SO₂R⁹, SO₂NR⁹R¹⁰, OH, C1 to 6 alkyl, C1 to 6 alkoxy, CN, CONR²R³, NR²COR³ and COR³, said alkoxy being optionally further substituted by C1 to 6 alkoxy, and said alkyl
25 being optionally further substituted by hydroxy or C1 to 6 alkoxy.

- In yet another embodiment, R¹ represents an aromatic ring system selected from phenyl, biphenyl, naphthyl or a five- or six-membered heteroaromatic ring containing 1 to 3 heteroatoms independently selected from O, N and S, said aromatic ring being optionally
30 substituted by one or more substituents independently selected from halogen, C1 to 6 alkyl and C1 to 6 alkoxy, said alkoxy being optionally further substituted by C1 to 6 alkoxy.

In yet another embodiment, R^1 represents an optionally substituted phenyl.

In yet another embodiment, R^1 represents an optionally substituted pyridyl.

5

In yet another embodiment, L represents C1 to 7 alkylene and R^1 represents H.

10

In yet another embodiment, L represents an optionally substituted C1 to 3 alkylene and R^1 represents a saturated or partially unsaturated 3- to 7-membered ring optionally incorporating one or two heteroatoms selected independently from O, N and S, and optionally incorporating a carbonyl group, said ring being optionally substituted by one or more substituents independently selected from halogen, SO_2R^9 , $SO_2NR^9R^{10}$, OH, C1 to 6 alkyl, C1 to 6 alkoxy, CN, $CONR^2R^3$, NR^2COR^3 and COR^3 , said alkoxy being optionally further substituted by C1 to 6 alkoxy, and said alkyl being optionally further substituted by hydroxy or C1 to 6 alkoxy.

15

20

In yet another embodiment, L represents an optionally substituted C1 to 3 alkylene and R^1 represents a saturated or partially unsaturated 3- to 7-membered ring optionally incorporating one or two heteroatoms selected independently from O, N and S, and optionally incorporating a carbonyl group, said ring being optionally substituted by one or more substituents independently selected from halogen, C1 to 6 alkyl and C1 to 6 alkoxy, said alkoxy being optionally further substituted by C1 to 6 alkoxy.

25

30

In yet another embodiment, L represents optionally substituted C1 to 3 alkylene and R^1 represents an aromatic ring system selected from phenyl, biphenyl, naphthyl or a five- or six-membered heteroaromatic ring containing 1 to 3 heteroatoms independently selected from O, N and S; said aromatic ring being optionally substituted by one or more substituents independently selected from halogen, SO_2R^9 , $SO_2NR^9R^{10}$, OH, C1 to 6 alkyl, C1 to 6 alkoxy, CN, $CONR^2R^3$, NR^2COR^3 and COR^3 , said alkoxy being optionally further substituted by C1 to 6 alkoxy, and said alkyl being optionally further substituted by hydroxy or C1 to 6 alkoxy.

In yet another embodiment, L represents optionally substituted C1 to 3 alkylene and R¹ represents an aromatic ring system selected from phenyl, biphenyl, naphthyl or a five- or six-membered heteroaromatic ring containing 1 to 3 heteroatoms independently selected from O, N and S, said aromatic ring being optionally substituted by one or more substituents independently selected from halogen, C1 to 6 alkyl and C1 to 6 alkoxy, said alkoxy being optionally further substituted by C1 to 6 alkoxy.

In yet another embodiment, X represents S, Y represents O, L represents optionally substituted C1 to 3 alkylene and R¹ represents optionally substituted phenyl.

In yet another embodiment, X represents S, Y represents O, L represents optionally substituted C1 to 3 alkylene and R¹ represents optionally substituted pyridyl.

In yet another embodiment, X represents S, Y represents O, L represents C1 to 3 alkylene, substituted with C1 to 6 alkoxy and R¹ represents hydrogen.

Particular compounds of the invention include:

1-butyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-isobutyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-(pyridin-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-(2-fluoro-benzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-[2-(2-methoxyethoxy)-3-propoxybenzyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-(6-ethoxy-pyridin-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-piperidin-3-ylmethyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-butyl-4-thioxo-1,3,4,5-tetrahydro-2H-pyrrolo[3,2-d]pyrimidin-2-one;

1-(2-isopropoxyethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-(2-methoxy-2-methylpropyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-(2-ethoxy-2-methylpropyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

1-(piperidin-4-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(1-methylpiperidin-3-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[2-hydroxy-2-(4-methoxyphenyl)ethyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
5 d]pyrimidin-4-one;
1-(2-methoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(3-methoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(2,4-dimethoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(3-chloropyridin-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
10 one;
1-{[3-(2-ethoxyethoxy)pyridin-2-yl]methyl}-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(6-oxo-1,6-dihydropyridin-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
15 1-(1H-indol-3-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(1H-benzimidazol-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(5-chloro-1H-indol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
20 1-[(5-fluoro-1H-indol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(1H-indol-6-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(1H-indol-5-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(5-fluoro-1H-indol-3-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
25 one;
1-(1H-imidazol-5-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(1H-imidazol-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(5-chloro-1H-benzimidazol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
30 1-[(4,5-dimethyl-1H-benzimidazol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
7-bromo-1-isobutyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one; and

1-(3-chlorophenyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one.

and pharmaceutically acceptable salts thereof.

A further aspect of the invention is the use of the novel compounds of formula (I) as a
5 medicament.

A further aspect of the invention is the use of a compound of formula (I), or a
pharmaceutically acceptable salt thereof, in the manufacture of a medicament, for the
treatment or prophylaxis of diseases or conditions in which inhibition of the enzyme MPO
10 is beneficial.

A further aspect of the invention provides the use of a compound of formula (I), or a
pharmaceutically acceptable salt thereof, in the manufacture of a medicament, for the
treatment or prophylaxis of neuroinflammatory disorders, cardio- and cerebrovascular
15 atherosclerotic disorders and peripheral artery disease and respiratory disorders such as
chronic obstructive pulmonary disease.

Another further aspect of the invention provides the use of a compound of formula (I), or a
pharmaceutically acceptable salt thereof, in the manufacture of a medicament, for the
20 treatment or prophylaxis of multiple sclerosis. Treatment may include slowing progression
of disease.

Another further aspect of the present invention provides the use of a compound of formula
(I) or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament, for
25 the treatment or prophylaxis of atherosclerosis by preventing and/or reducing the formation
of new atherosclerotic lesions or plaques and/or by preventing or slowing progression of
existing lesions and plaques.

Another further aspect of the present invention provides the use of a compound of formula
30 (I) or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament, for

the treatment or prophylaxis of atherosclerosis by changing the composition of the plaques to reduce the risk of plaque rupture and atherothrombotic events.

Another further aspect of the present invention provides the use of a compound of formula
5 (I) or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament, for the treatment or prophylaxis of respiratory disorders, such as chronic obstructive pulmonary disease. Treatment may include slowing progression of disease.

According to the invention, there is also provided a method of treating, or reducing the risk
10 of, diseases or conditions in which inhibition of the enzyme MPO is beneficial which comprises administering to a person suffering from or at risk of, said disease or condition, a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

15 Further, there is also provided a method of treating, or reducing the risk of, neuroinflammatory disorders, cardio- and cerebrovascular atherosclerotic disorders or peripheral artery disease, or respiratory disorders, such as chronic obstructive pulmonary disease, in a person suffering from or at risk of, said disease or condition, wherein the method comprises administering to the person a therapeutically effective amount of a
20 compound of formula (I), or a pharmaceutically acceptable salt thereof.

Further, there is also provided a method of treating, or reducing the risk of, multiple sclerosis in a person suffering from or at risk of, said disease or condition, wherein the method comprises administering to the person a therapeutically effective amount of a
25 compound of formula (I), or a pharmaceutically acceptable salt thereof.

There is also provided a method of treating, or reducing the risk of atherosclerosis by preventing and/or reducing the formation of new atherosclerotic lesions or plaques and /or by preventing or slowing progression of existing lesions and plaques in a person suffering
30 from or at risk of, said disease or condition, wherein the method comprises administering

to the person a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

There is also provided a method of treating, or reducing the risk of atherosclerosis by
5 changing the composition of the plaques so as to reduce the risk of plaque rupture and atherothrombotic events in a person suffering from or at risk of, said disease or condition, wherein the method comprises administering to the person a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

10 In another aspect the invention provides a pharmaceutical formulation comprising a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier, for use in the treatment or prophylaxis of diseases or conditions in which inhibition of the enzyme MPO is beneficial.

15 In a further aspect the invention provides a pharmaceutical formulation comprising a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier, for use in the treatment or prophylaxis of neuroinflammatory disorders.


20 In a further aspect the invention provides a pharmaceutical formulation comprising a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier, for use in the treatment or prophylaxis of multiple sclerosis, cardio- and
25 cerebrovascular atherosclerotic disorders and peripheral artery disease and respiratory disorders, such as chronic obstructive pulmonary disease.

In another aspect the present invention provides a pharmaceutical formulation comprising a therapeutically effective amount of a compound of formula (I), or a pharmaceutically
30 acceptable salt thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent

or carrier, for use in the treatment or prophylaxis of atherosclerosis by preventing and reducing the formation of new atherosclerotic lesions and/or plaques and/or by preventing or slowing progression of existing lesions and plaques.

5 In another aspect the present invention provides a pharmaceutical formulation comprising a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier, for use in the treatment or prophylaxis of atherosclerosis by changing the composition of the plaques so as to reduce the risk of plaque rupture and atherothrombotic
10 events.

According to the invention, there is further provided a process for the preparation of the novel compounds of formula (I), or a pharmaceutically acceptable salt, tautomer, enantiomer, diastereomer or racemate thereof which comprises reaction of a compound of
15 formula (II),




wherein R¹, R¹² and L are as defined in formula (I) and R represents C1 to 6 alkoxy with
20 the oxygen in a direct bond to the carbonyl in formula (II) with alkoxy as defined above or NH₂;

with a C1 to 6 alkoxycarbonyl isothiocyanate or with a phenylcarbonyl isothiocyanate, wherein the phenyl group is optionally substituted by one or more groups selected independently from C1 to 6 alkyl, halogen, C1 to 6 alkoxy, NO₂, OH, CN, C1 to 6
25 alkylamino or NH₂; and where necessary converting the resultant compound of formula (I), or another salt thereof, into a pharmaceutically acceptable salt thereof; or converting the


resultant compound of formula (I) into a further compound of formula (I); and where desired converting the resultant compound of formula (I) into an optical isomer thereof.

In the process, a compound of formula (II) and the alkoxycarbonyl isothiocyanate or the phenylcarbonyl isothiocyanate are dissolved or suspended in a suitable dry organic solvent such as dichloromethane or and stirred at 0 to 30 degrees for example at ambient temperature until reaction is complete, typically for between 5 to 60 minutes, but if necessary, overnight. Preferably the alkoxycarbonyl isothiocyanate is ethoxycarbonyl isothiocyanate and the phenylcarbonyl isothiocyanate is preferably benzoyl isothiocyanate. Following a standard work-up the intermediate product is then optionally purified before treatment with a base, such as sodium ethoxide in ethanol, aqueous sodium hydroxide or ammonia in solution, ammonia in methanol, to give the required compound of formula (I). The cyclization is carried out at an elevated temperature either in an oil bath or in a microwave reactor. See, for example, Norman et al, *J. Med. Chem.* 2000, 43, 4288-4312.

When ammonia in methanol is used, a pressure vessel is preferably used.




Compounds of formula (II) may be prepared by reaction of a compound of formula (III), wherein R^{12} is as defined in Formula I, R is as defined in formula II (see for example Furneaux et al, *J. Org. Chem.* 1999, 64, 8411-8412), and may be carried out by




a) reductive amination. In the process, a compound of formula (III) may be mixed with an aldehyde of formula (IV), wherein R^1 is defined as in formula I, in the presence of a reducing agent such as sodium cyanoborohydride or sodium triacetoxyborohydride. An

acid, preferably acetic acid, may be added to catalyze the reaction. The reaction may be performed in a solvent such as methanol between ambient temperature and 50 °C, preferably at ambient temperature. Following a standard work-up the product is then optionally purified by flash column chromatography. See for example Suzuki et al, *Chem. Pharm. Bull.* 2002, 50, 1163-1168, or Furneaux, R.H., Tyler, P.C., *J. Org. Chem.* 1999, 64, 8411-8412.



- b) alkylation. In the process, a mesylate of formula (V), wherein R^1 is defined as above, may be added to a stirred solution of a compound of formula (III), potassium iodide and a base, preferably potassium carbonate. The reaction may be performed in a solvent, such as N,N-dimethylformamide, at an elevated reaction temperature, preferably at 85 °C. The reaction mixture may be worked up by extraction and then purified by flash column chromatography to give a compound of formula (II).



- c) cross-coupling of a compound of formula (III) with a suitable aryl of a compound (VI), wherein R^1 is defined as above and $Halo$ is halogen, preferably bromo, to give a compound of formula (II). The reaction may be carried out using a suitable palladium catalyst such as $Pd_2(dba)_3$ or $Pd(OAc)_2$ together with a suitable ligand such as BINAP. A suitable base, such as cesium carbonate, may be used in the reaction in a suitable solvent such as tetrahydrofuran, dioxane or toluene, which is performed in the temperature range between 80 °C and 100 °C. See for example, J.P. Wolfe, S.L. Buchwald *J. Org. Chem.* 2000, 65, 1144-1157.

Compounds of formula (II) are either known in the literature or may be prepared using known methods that will be readily apparent to the man skilled in the art. See, for example, Suzuki et al, *Chem. Pharm. Bull.* 2002, 50, 1163-1168, or Furneaux, R.H., Tyler, P.C., *J. Org. Chem.* 1999, 64, 8411-8412.

Compounds of formula (IV), (V) and (VI) are either commercially available or may be prepared using methods that are well-known in the literature.

The present invention includes compounds of formula (I) in the form of salts. Suitable salts include those formed with organic or inorganic acids or organic or inorganic bases. Such salts will normally be pharmaceutically acceptable although salts of non-pharmaceutically acceptable acids or bases may be of utility in the preparation and purification of the compound in question. Thus, preferred acid addition salts include those formed from hydrochloric, hydrobromic, sulphuric, phosphoric, citric, tartaric, lactic, pyruvic, acetic, succinic, fumaric, maleic, methanesulphonic and benzenesulphonic acids. Preferred base addition salts include those in which the cation is sodium, potassium, calcium, aluminium, lithium, magnesium, zinc, choline, ethanolamine or diethylamine.

Salts of compounds of formula (I) may be formed by reacting the compound, or a salt, enantiomer or racemate thereof, with one or more equivalents of the appropriate acid or base. The reaction may be carried out in a solvent or medium in which the salt is insoluble or in a solvent in which the salt is soluble, for example, water, dioxan, ethanol, tetrahydrofuran or diethyl ether, or a mixture of solvents, which may be removed *in vacuo* or by freeze drying. The reaction may also be a metathetical process or it may be carried out on an ion exchange resin.

The compounds of the invention and intermediates thereto may be isolated from their reaction mixtures and, if necessary further purified, by using standard techniques.

The compounds of formula (I) may exist in enantiomeric forms. Therefore, all enantiomers, diastereomers, racemates, tautomers and mixtures thereof are included within the scope of the invention. The various optical isomers may be isolated by separation of a racemic mixture of the compounds using conventional techniques, for example, fractional crystallisation, or
5 HPLC. Alternatively, the various optical isomers may be prepared directly using optically active starting materials.

Intermediate compounds may also exist in enantiomeric forms and may be used as purified enantiomers, diastereomers, racemates or mixtures.

10 The compounds of formula (I) may exist in tautomeric forms. All such tautomers and mixtures of tautomers are included within the scope of the invention.

Intermediate compounds may also exist in tautomeric forms and may be used as purified
15 tautomers or mixtures.

The compounds of formula (I) and their pharmaceutically acceptable salts are useful because they possess pharmacological activity as inhibitors of the enzyme MPO.

20 The compounds of formula (I) and their pharmaceutically acceptable salts are indicated for use in the treatment or prophylaxis of diseases or conditions in which modulation of the activity of the enzyme myeloperoxidase (MPO) is desirable. In particular, linkage of MPO activity to disease has been implicated in neuroinflammatory diseases. Therefore the compounds of the present invention are particularly indicated for use in the treatment of
25 neuroinflammatory conditions or disorders in mammals including man. The compounds are also indicated to be useful in the treatment of cardio- and cerebrovascular atherosclerotic disorders or peripheral artery disease. The compounds are also indicated to be useful in the treatment of respiratory disorders, such as disorders of the respiratory tract: obstructive diseases of the airways including: asthma, including bronchial, allergic, intrinsic, extrinsic,
30 exercise-induced, drug-induced (including aspirin and NSAID-induced) and dust-induced

asthma, both intermittent and persistent and of all severities, and other causes of airway hyper-responsiveness; chronic obstructive pulmonary disease (COPD); bronchitis, including infectious and eosinophilic bronchitis; emphysema; bronchiectasis; cystic fibrosis; sarcoidosis; farmer's lung and related diseases; hypersensitivity pneumonitis; lung
5 fibrosis, including cryptogenic fibrosing alveolitis, idiopathic interstitial pneumonias, fibrosis complicating anti-neoplastic therapy and chronic infection, including tuberculosis and aspergillosis and other fungal infections; complications of lung transplantation; vasculitic and thrombotic disorders of the lung vasculature, and pulmonary hypertension; antitussive activity including treatment of chronic cough associated with inflammatory and
10 secretory conditions of the airways, and iatrogenic cough; acute and chronic rhinitis including rhinitis medicamentosa, and vasomotor rhinitis; perennial and seasonal allergic rhinitis including rhinitis nervosa (hay fever); nasal polyposis; acute viral infection including the common cold, and infection due to respiratory syncytial virus, influenza, coronavirus (including SARS) and adenovirus. Such conditions or disorders will be readily
15 apparent to the man skilled in the art.

Conditions or disorders that may be specifically mentioned include multiple sclerosis, Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis and stroke, as well as other inflammatory diseases or conditions such as asthma, chronic obstructive
20 pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, acute respiratory distress syndrome, sinusitis, rhinitis, psoriasis, dermatitis, uveitis, gingivitis, atherosclerosis, myocardial infarction, stroke, coronary heart disease, ischaemic heart disease, restenosis, inflammatory bowel disease, renal glomerular damage, liver fibrosis, sepsis, proctitis, rheumatoid arthritis, and inflammation associated with reperfusion injury, spinal cord
25 injury and tissue damage/scarring/adhesion/rejection. Lung cancer has also been suggested to be associated with high MPO levels. The compounds are also expected to be useful in the treatment of pain.

Prophylaxis is expected to be particularly relevant to the treatment of persons who have
30 suffered a previous episode of, or are otherwise considered to be at increased risk of, the

disease or condition in question. Persons at risk of developing a particular disease or condition generally include those having a family history of the disease or condition, or those who have been identified by genetic testing or screening to be particularly susceptible to developing the disease or condition.

5

For the above-mentioned therapeutic indications, the dosage administered will, of course, vary with the compound employed, the mode of administration and the treatment desired. However, in general, satisfactory results are obtained when the compounds are administered at a dosage of the solid form of between 1 mg and 2000 mg per day.

10

The compounds of formulae (I), and pharmaceutically acceptable derivatives thereof, may be used on their own, or in the form of appropriate pharmaceutical compositions in which the compound or derivative is in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier. Thus, another aspect of the invention concerns a pharmaceutical composition comprising a novel compound of formula (I), or a pharmaceutically acceptable salt thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier. Administration may be by, but is not limited to, enteral (including oral, sublingual or rectal), intranasal, inhalation, intravenous, topical or other parenteral routes. Conventional procedures for the selection and preparation of suitable pharmaceutical formulations are described in, for example, "Pharmaceuticals - The Science of Dosage Form Designs", M. E. Aulton, Churchill Livingstone, 1988. The pharmaceutical composition preferably comprises less than 80% and more preferably less than 50% of a compound of formulae (I), or a pharmaceutically acceptable salt thereof.

15

20

25

There is also provided a process for the preparation of such a pharmaceutical composition which comprises mixing the ingredients.

30

Examples of pharmaceutical composition

The following illustrate representative pharmaceutical dosage forms containing a compound of formula I, or salts, solvates or solvated salts thereof, (hereafter compound X), for preventive or therapeutic use in mammals:

5

(a): Tablet	mg/tablet
Compound X	100
Lactose	182.75
Croscarmellose sodium	12.0
Maize starch paste (5% w/v paste)	2.25
Magnesium stearate	3.0

(b): Capsule	mg/capsule
Compound X	10
Lactose	488.5
Magnesium stearate	1.5

(c): Injection	(50 mg/ml)
Compound X	5.0% w/v
1M Sodium hydroxide solution	15.0% v/v
0.1M Hydrochloric acid	(to adjust pH to 7.6)
Polyethylene glycol 400	4.5% w/v
Water for injection	up to 100%

The above compositions may be obtained by conventional procedures well known in the pharmaceutical art.

10

The invention further relates to combination therapies wherein a compound of formula (I) or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition or formulation comprising a compound of formula (I), is administered concurrently or

sequentially with therapy and/or an agent for the treatment of any one of cardio- and cerebrovascular atherosclerotic disorders and peripheral artery disease.

In particular, a compound of formula (I) or a pharmaceutically acceptable salt thereof may be administered in association with compounds from one or more of the following groups:

5 1) anti-inflammatory agents, for example

a) NSAIDs (e.g. acetylsalicylic acid, ibuprofen, naproxen, flurbiprofen, diclofenac, indometacin);

b) leukotriene synthesis inhibitors (5-LO inhibitors e.g. AZD4407, Zileuton, licofelone, CJ13610, CJ13454; FLAP inhibitors e.g. BAY-Y-1015, DG-031, MK591, MK886, A81834; LTA4 hydrolase inhibitors e.g. SC56938, SC57461A);

c) leukotriene receptor antagonists (e.g. CP195543, amelubant, LY293111, accolat, MK571);

2) anti-hypertensive agents, for example

15 a) beta-blockers (e.g. metoprolol, atenolol, sotalol);

b) angiotensin converting enzyme inhibitors (e.g. captopril, ramipril, quinapril, enalapril);

c) calcium channel blockers (e.g. verapamil, diltiazem, felodipine, amlodipine);

d) angiotensin II receptor antagonists (e.g. irbesartan, candesartan, telmisartan, losartan);

3) anti-coagulantia, for example

a) thrombin inhibitors (e.g. ximelagatran), heparines, factor Xa inhibitors;

b) platelet aggregation inhibitors (e.g. clopidogrel, ticlopidine, prasugel, AZ4160);

4) modulators of lipid metabolism, for example

25 a) insulin sensitizers such as PPAR agonists (e.g. pioglitazone, rosiglitazone, Galida, muraglitazar, gefemrozil, fenofibrate);

b) HMG-CoA reductase inhibitors, statins (e.g. simvastatin, pravastatin, atorvastatin, rosuvastatin, fluvastatin);

c) cholesterol absorption inhibitors (e.g. ezetimibe);

30 d) IBAT inhibitors (e.g. AZD-7806);

e) LXR agonists (e.g. GW-683965A, T-0901317);

f) FXR receptor modulators;

- g) phospholipase inhibitors;
- 5) anti-anginal agents, for example, nitrates and nitrites;
- 6) modulators of oxidative stress, for example, anti-oxidants (probutol).

General Methods

All solvents used were analytical grade and commercially available anhydrous solvents were routinely used for reactions. Reactions were typically run under an inert atmosphere of nitrogen or argon.

^1H and ^{13}C NMR spectra were recorded at 400 MHz for proton and 100 MHz for carbon-13 either on a Varian Unity+ 400 NMR Spectrometer equipped with a 5mm BBO probe head with Z-gradients, or a Bruker Avance 400 NMR spectrometer equipped with a 60 μl dual inverse flow probe head with Z-gradients, or a Bruker DPX400 NMR spectrometer equipped with a 4-nucleus probe head equipped with Z-gradients. Unless specifically noted in the examples, spectra were recorded at 400 MHz for proton and 100 MHz for carbon-13. The following reference signals were used: the middle line of DMSO- d_6 δ 2.50 (^1H), δ 39.51 (^{13}C); the middle line of CD_3OD δ 3.31 (^1H) or δ 49.15 (^{13}C); acetone- d_6 2.04 (^1H), 206.5 (^{13}C); and CDCl_3 δ 7.26 (^1H), the middle line of CDCl_3 δ 77.16 (^{13}C) (unless otherwise indicated).

Mass spectra were recorded on a Waters LCMS consisting of an Alliance 2795 (LC), Waters PDA 2996, and ELS detector (Sedex 75) and a ZMD single quadrupole mass spectrometer. The mass spectrometer was equipped with an electrospray ion source (ES) operated in a positive or negative ion mode. The capillary voltage was 3 kV and cone voltage was 30 V. The mass spectrometer was scanned between m/z 100-600 with a scan time of 0.7s. The column temperature was set to 40 $^\circ\text{C}$. The Diode Array Detector was scanned from 200-400 nm. The temperature of the ELS detector was adjusted to 40 $^\circ\text{C}$ and the pressure was set to 1.9 bar. For LC separation a linear gradient was applied starting at 100% A (A: 10 mM NH_4OAc in 5% MeCN) and ending at 100% B (B: MeCN) after four

minutes. The column used was a X-Terra MS C8, 3.0 x 50; 3.5 μm (Waters) run at 1.0 mL/min.

Alternatively, mass spectra was performed on a GC-MS (GC 6890, 5973N MSD) supplied
5 by Agilent Technologies. The column used was a VF-5 MS, ID 0.25 mm x 30m, 0.25 μm (Varian Inc.). A linear temperature gradient was applied starting at 40 °C (hold 1 min) and ending at 300 °C (hold 1 min), 25 °C/minute. The MS was equipped with a CI ion source and the reactant gas was methane. The MS was scanned between m/z 50-500 and the scan speed was set to 3.25 scan/s. The MS was equipped with an EI ion source. The MS was
10 scanned between m/z 50-500 and the scan speed was set to 3.25 scan/s. The electron voltage was set to 70 eV.

HPLC analyses were performed on an Agilent HP1100 system consisting of G1379A Micro Vacuum Degasser, G1312A Binary Pump, G1367A Well plate auto-sampler,
15 G1316A Thermostatted Column Compartment and G1315B Diode Array Detector. Column: X-Terra MS, Waters, 3.0 x 100 mm, 3.5 μm . The column temperature was set to 40 °C and the flow rate to 1.0 ml/min. The Diode Array Detector was scanned from 210-300 nm, step and peak width were set to 2 nm and 0.05 min, respectively. A linear gradient was applied, starting at 100 % A (A: 10 mM NH_4OAc in 5 % MeCN) and ending at 100%
20 B (B: MeCN), in 6 min.

Microwave heating was performed in an Initiator or Smith Synthesizer Single-mode microwave cavity producing continuous irradiation at 2450 MHz.

25 A typical workup procedure after a reaction consisted of extraction of the product with a solvent such as ethyl acetate, washing with water followed by drying of the organic phase over MgSO_4 or Na_2SO_4 filtration and concentration of the solution *in vacuo*.

Thin layer chromatography (TLC) was performed on Merck TLC-plates (Silica gel 60 F₂₅₄)
30 and UV visualized the spots. Flash column chromatography was performed on a Combi Flash[®] Companion[™] using RediSep[™] normal-phase flash columns. Typical solvents used

for flash column chromatography were mixtures of chloroform/methanol, dichloromethane/methanol and heptane/ethyl acetate.

Preparative chromatography was run on a Waters autopurification HPLC with a diode array detector. Column: XTerra MS C8, 19 x 300 mm, 10 μ m. Narrow gradients with MeCN/(95:5 0.1M NH₄OAc:MeCN) were used at a flow rate of 20 ml/min. Alternatively, another column was used; Atlantis C18 19 x 100 mm, 5 μ m column. Gradient with acetonitrile/0.1M ammonium acetate in 5% acetonitrile in MilliQ Water, run from 0% to 35-50% acetonitrile, in 15 min. Flow rate: 15 ml/min. Alternatively, purification was achieved on a semi preparative Shimadzu LC-8A HPLC with a Shimadzu SPD-10A UV-vis.-detector equipped with a Waters Symmetry[®] column (C18, 5 μ m, 100 mm x 19 mm). Narrow gradients with MeCN/0.1% trifluoroacetic acid in MilliQ Water were used at a flow rate of 10 ml/min.

Recrystallization was typically performed in solvents or solvent mixtures such as ether, ethyl acetate/heptane and methanol/water.

The following abbreviations have been used:

aq. aqueous;

BINAP 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl

equiv. equivalent;

DMF N,N-dimethylformamide;

DMSO dimethylsulfoxide;

DIBAL diisobutylaluminium hydride;

Et₃N triethyl amine;

HOAc acetic acid;

NaBH₄ sodium borohydride;

NaCNBH₃ sodium cyanoborohydride;

Pd₂(dba)₃ tris(dibenzylideneacetone)dipalladium;

Pd(OAc)₂ palladium diacetate;

r.t. room temperature;

TBDMSCl tert-butyldimethylsilyl chloride;

TEMPO 2,2,6,6-tetramethyl-1-piperidinyloxy


THF tetrahydrofuran.

Starting materials used were either available from commercial sources or prepared

according to literature procedures and had experimental data in accordance with those reported. The following is an example of a starting material that was prepared:


3-Amino-1H-pyrrole-2-carboxylic acid ester: Furneaux, R.H., Tyler, P.C., *J. Org. Chem.* 1999, 64, 8411-8412.

General Method A



A reaction mixture of the amino pyrrole ester **A1** (1.0 equiv.), the aldehyde **A2** (1.0 to 2.0 equiv.) and NaCNBH₃ (1.0 equiv.) in methanol was stirred at r.t. for 24 h. In some examples, acetic acid (1 to 2 equiv.) was added to catalyze the reaction. If the reaction was not complete after 24 h (monitored by TLC or LC-MS), more aldehyde **A2** was added and the mixture was stirred at r.t. until the reaction was complete. The mixture was then evaporated onto silica gel and purified by flash column chromatography.

General Method B



Ethoxycarbonyl isothiocyanate **B2** (1.0 to 1.2 equiv.) was added to the amino pyrrole ester **B1** (1.0 equiv.) in CH₂Cl₂ and the mixture was stirred at r.t. for 5 to 60 minutes, or overnight. Water was added and the aqueous phase was extracted with CH₂Cl₂. The organic phase was combined, dried (MgSO₄) and concentrated. The crude ring opened intermediate was purified by flash column chromatography. The intermediate product was dissolved in 1M NaOEt in EtOH (1.1-1.5 equiv.) and heated in a microwave reactor for 10 minutes at 120 °C. The pH was adjusted to neutral pH with 2M HCl; the solid was collected by filtration and washed with water. The crude product was purified using preparative HPLC, or by flash column chromatography or by recrystallization.

The invention is illustrated, but in no way limited, by the following examples. Except where otherwise indicated, the compounds of Examples 1a to 4a and 5c and 7b were prepared using the procedure of General Method A, and the compounds of Examples 1b to 4b and 5d and 7c were prepared using the procedure of General Method B.

Example 1

1-Butyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 3-(Butylamino)-1H-pyrrole-2-carboxylic acid ethyl ester

The title compound was obtained as an oil in 60% yield starting from 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.81 g, 5.26 mmol) and butyraldehyde (0.47 + 0.55 mL, 11.4 mmol).

¹H NMR (DMSO-d₆) δ ppm 10.71 (1H, br s), 6.74 (1H, t, J=3.1 Hz), 5.62 (1H, t, J=2.6 Hz), 5.19 (1H, s), 4.17 (2H, q, J=7.0 Hz), 3.04 (2H, q, J=6.6 Hz), 1.50 (2H, m), 1.34 (2H, m), 1.25 (3H, t, J=7.0 Hz), 0.90 (3H, t, J=7.3 Hz);

MS (ESI) m/z 211 ($M+1$).

(b) 1-Butyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound was obtained as a solid in 44% yield starting from 3-(butylamino)-1H-pyrrole-2-carboxylic acid ethyl ester (0.10 g, 0.48 mmol) and ethoxycarbonyl isothiocyanate (0.06 mL, 0.58 mmol).

^1H NMR (DMSO- d_6) δ ppm 12.38 (1H, s), 12.10 (1H, s), 7.37 (1H, d, $J=2.9$), 6.31 (1H, d, $J=2.6$ Hz), 4.36 (2H, m), 1.69 (2H, m), 1.38 (2H, m), 0.92 (3H, t, $J=7.5$ Hz);

MS (ESI) m/z 224 ($M+1$).

Example 2

1-Isobutyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 3-(Isobutylamino)-1H-pyrrole-2-carboxylic acid ethyl ester

The title compound was obtained as an oil in 71% yield starting from 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.40 g, 2.59 mmol) and isobutyraldehyde (0.26 + 0.07 mL, 3.61 mmol).

^1H NMR (DMSO- d_6) δ ppm 10.68 (1H, s), 6.74 (1H, t, $J=3.0$ Hz), 5.62 (1H, t, $J=2.4$ Hz), 5.30 (1H, br s), 4.18 (2H, q, $J=7.2$ Hz), 2.88 (2H, t, $J=6.4$ Hz), 1.79 (1H, m), 1.26 (3H, t, $J=7.1$ Hz), 0.90 (3H, s), 0.89 (3H, s);

^{13}C NMR (DMSO- d_6) δ ppm 160.9, 124.2, 95.0, 58.2, 52.4, 27.9, 20.0, 14.7;

MS (ESI) m/z 211 ($M+1$).

(b) 1-Isobutyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound was obtained as a solid in 24% yield starting from 3-(isobutylamino)-1H-pyrrole-2-carboxylic acid ethyl ester (0.38 g, 1.79 mmol) and ethoxycarbonyl isothiocyanate (0.24 mL, 2.15 mmol).

^1H NMR (DMSO- d_6) δ ppm 12.36 (1H, br s), 12.13 (1H, br s), 7.35 (1H, d, $J=2.8$ Hz), 6.34 (1H, d, $J=2.8$ Hz), 4.21 (2H, d, $J=7.33$ Hz), 2.44 (1H, m), 0.91 (3H, s), 0.90 (3H, s);

^{13}C NMR (DMSO- d_6) δ ppm 172.8, 152.4, 137.3, 127.7, 113.6, 97.1, 56.2, 26.4, 19.7;

MS (ESI) m/z 224 ($M+1$).

Example 3

1-(Pyridin-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 3-[(Pyridin-2-ylmethyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester

The title compound was obtained as an oil in 54% yield starting from 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.40 g, 2.59 mmol) and 2-pyridinecarboxaldehyde (0.27 + 0.07 mL, 3.55 mmol).

¹H NMR (DMSO-d₆) δ ppm 10.77 (1H, br s), 8.52 (1H, d, J=4.0 Hz), 7.80-7.67 (1H, m), 7.35 (1H, d, J=7.8 Hz), 7.25 (1H, dd, J=7.3, 5.0 Hz), 6.71 (1H, t, J=3.0 Hz), 6.10 (1H, br s), 5.57 (1H, t, J=2.4 Hz), 4.37 (2H, d, J=5.8 Hz), 4.21 (2H, q, J=7.2 Hz), 1.29 (3H, t, J=7.1 Hz);

MS (ESI) m/z 246 (M +1).

(b) 1-(Pyridin-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound was obtained as a solid in 14% yield starting from 3-[(pyridin-2-ylmethyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester (0.34 g, 1.39 mmol) and ethoxycarbonyl isothiocyanate (0.19 mL, 1.66 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.34 (2H, br s), 8.49 (1H, d, J=4.5 Hz), 7.73 (1H, m), 7.29 (1H, d, J=2.8 Hz), 7.27 (1H, m), 7.21 (1H, d, J=7.8 Hz), 6.09 (1H, d, J=2.8 Hz), 5.75 (2H, s);

¹³C NMR (DMSO-d₆) δ ppm 173.4, 155.2, 152.6, 149.1, 137.1, 136.8, 127.9, 122.4, 121.2, 113.6, 96.9, 54.2;

MS (ESI) m/z 259 (M +1).

Example 4

1-(2-Fluoro-benzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 3-(2-Fluoro-benzylamino)-1H-pyrrole-2-carboxylic acid ethyl ester

The title compound was obtained as an oil in quantitative yield starting from 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.50 g, 3.2 mmol), and 2-fluorobenzaldehyde (0.34 mL, 3.2 mmol) using the general procedure A but with the following modifications. After 5 h more NaCNBH₃ (100 mg, 1.6 mmol) was added followed by more 2-fluorobenzaldehyde (120 mg, 1 mmol), and the reaction was then stirred overnight.

¹H NMR (DMSO-d₆) δ ppm 10.76 (1H, br s), 7.47 (1H, m), 7.38 (1H, m), 7.28 (2H, m), 6.70 (1H, m), 5.74 (1H, br s), 5.61 (1H, m), 4.34 (2H, m), 4.18 (2H, q, J=7.1 Hz), 1.25 (3H, t, J=7.1 Hz);

MS (ES) m/z 263 (M +1).

(b) *1-(2-Fluoro-benzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one*

The title compound was obtained as a solid in 45% yield starting from 3-(2-fluoro-benzylamino)-1H-pyrrole-2-carboxylic acid ethyl ester (0.85 g, 3.2 mmol) and ethoxycarbonyl isothiocyanate (0.44 mL, 3.9 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.41 (2H, br s.), 7.33 (2H, m), 7.24 (1H, m), 7.10 (1H, m, J=7.5, 7.5 Hz), 7.01 (1H, m, J=7.1 Hz, 7.1 Hz), 6.12 (1H, d, J=2.8 Hz), 5.72 (2H, s);
¹³C NMR (DMSO-d₆) δ 173.9, 161.4, 159.0, 152.9, 137.1, 129.6, 129.5, 128.5, 128.1, 128.1, 125.0, 124.9, 123.2, 123.0, 115.8, 115.6, 114.1, 96.9, 47.1, 47.1;
MS (ESI) m/z 276 (M +1).

Example 5

1-[2-(2-Methoxyethoxy)-3-propoxybenzyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) *3-Hydroxy-2-(2-methoxyethoxy)benzaldehyde*

2-Chloroethyl methyl ether (4.63 mL, 50.7 mmol) was added dropwise to a mixture of 2,3-dihydroxybenzaldehyde (7.0 g, 50.7 mmol), potassium iodide (8.41 g, 50.69 mmol) and potassium carbonate (7.71 g, 55.8 mmol) in DMF (80 mL). The resulting mixture was stirred at r.t. under a nitrogen atmosphere for two days and at 70 °C for two days. The reaction mixture was partitioned between saturated ammonium chloride (aq.) and CH₂Cl₂.

The water phase was re-extracted with CH₂Cl₂ and the combined organic phases were washed with brine, dried (Na₂SO₄) and evaporated onto silica. Purification by flash column chromatography (heptane/ethyl acetate gradient; 0 to 30% ethyl acetate) yielded a crude oil which was further purified by flash column chromatography (heptane/ethyl acetate gradient; 0 to 40% ethyl acetate) to yield the title compound (3.13 g, 31%) as an oil.

¹H NMR (DMSO-d₆) δ ppm 10.34 (1H, br s), 9.88 (1H, br s), 7.16 (2H, m), 7.05 (1H, m), 4.25 (2H, m), 3.60 (2H, m), 3.26 (3H, m);
¹³C NMR (DMSO-d₆) δ ppm 190.5, 150.7, 149.6, 129.9, 124.1, 122.6, 116.9, 72.1, 70.9, 57.9;
MS (ESI) m/z 197 (M +1).

(b) *2-(2-Methoxyethoxy)-3-propoxybenzaldehyde*

1-Iodopropane (3.09 mL, 31.60 mmol) was added to a solution of 3-hydroxy-2-(2-methoxyethoxy)benzaldehyde (3.1 g, 15.8 mmol) and potassium carbonate (4.37 g, 31.60

mmol) in DMF (80 mL) and the mixture was stirred at 100 °C overnight under a nitrogen atmosphere. The reaction mixture was partitioned between saturated ammonium chloride (aq.) and CH₂Cl₂. The organic phase was washed with brine, dried (Na₂SO₄) and concentrated to give the title compound in quantitative yield (3.8 g) as an oil. This material
5 was used in the next step without further purification.

¹H NMR (DMSO-d₆) δ ppm 10.37 (1H, s), 7.36 (1H, m, J=8.0 Hz), 7.26 (1H, m), 7.17 (1H, t, J=7.8 Hz), 4.28 (2H, m), 4.02 (2H, t, J=6.3 Hz), 3.62 (2H, m), 3.26 (3H, s), 1.80 (2H, m), 1.02 (3H, t, J=7.3 Hz);

¹³C NMR (DMSO-d₆) δ ppm 190.3, 152.1, 150.9, 129.6, 124.4, 119.4, 117.8, 72.5, 70.9,
10 70.0, 57.9, 22.1, 10.4;

MS (ESI) m/z 239 (M +1).

(c) 3-{[2-(2-Methoxyethoxy)-3-propoxybenzyl]amino}-1H-pyrrole-2-carboxylic acid ethyl ester

The title compound was obtained as an oil in quantitative yield starting from 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.35 g, 2.27 mmol) and 2-(2-methoxyethoxy)-3-propoxybenzaldehyde (0.47 + 0.08 g, 3.06 mmol).
15

¹H NMR (DMSO-d₆) δ ppm 10.69 (1H, br s), 7.13 (1H, s), 7.6.99 (2H, m), 6.70 (1H, m), 5.63 (1H, m), 4.92 (1H, t, J=5.7 Hz), 4.52 (2H, d, J=5.8 Hz), 4.19 (2H, m), 4.05 (2H, m), 3.92 (2H, t, J=6.4 Hz), 3.59 (2H, m), 3.32 (3H, s), 1.76 (2H, m), 1.26 (3H, t, J=7.1 Hz),
20 1.00 (3 H, m);

MS (ESI) m/z 377 (M +1).

(d) 1-[2-(2-Methoxyethoxy)-3-propoxybenzyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound was obtained as a solid in 13% yield starting from 3-{[2-(2-methoxyethoxy)-3-propoxybenzyl]amino}-1H-pyrrole-2-carboxylic acid ethyl ester (0.87
25 g, 2.31 mmol) and ethoxycarbonyl isothiocyanate (0.26 mL, 2.31 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.43 (1H, br s), 12.31 (1H, br s), 7.29 (1H, d, J=3.0 Hz), 6.96–6.85 (2H, m), 6.41 (1H, dd, J=7.3, 1.5 Hz), 6.02 (1H, d, J=2.8 Hz), 5.71 (2H, s), 4.23 (2H, m), 3.95 (2H, t, J=6.3 Hz), 3.65 (2H, m), 3.33 (3H, s), 1.82–1.72 (2H, m), 1.02 (3H, t, J=7.4 Hz);
30

MS (ESI) m/z 390 (M +1).

Example 6**1-(6-Ethoxy-pyridin-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one***(a) 6-Ethoxy-pyridine-2-carboxylic acid ethyl ester*

5 Ethyl iodide (2.3 mL, 28.8 mmol) was added to a suspension of 6-hydroxy-pyridine-2-carboxylic acid (1.0 g, 7.2 mmol) and silver(I) carbonate (4.0 g, 14.4 mmol) in CHCl₃ (70 mL). The suspension was stirred at ambient temperature for 3 days. Insoluble material was removed by filtration and the solid was washed with CHCl₃. The filtrate was concentrated to give the title product in quantitative yield (1.5 g) as an oil. This material was used in the
10 next step without further purification.

¹H NMR (CDCl₃) δ ppm 7.65 (2H, m), 6.88 (1H, m), 4.45 (2H, q, J=7.0 Hz), 4.41 (2H, q, J=7.3 Hz), 1.40 (6H, m);

MS (ESI) m/z 196 (M+1).

(b) (6-Ethoxy-pyridin-2-yl)-methanol

15 NaBH₄ (5.7 g, 151 mmol) was added in portions during 35 minutes to 6-ethoxy-pyridine-2-carboxylic acid ethyl ester (1.5 g, 7.5 mmol) in EtOH (75 mL). The resulting mixture was stirred at ambient temperature for two days. Water was added and the mixture was extracted with CH₂Cl₂. The organic phase was dried (Na₂SO₄), filtered and then concentrated to give the title product (0.85 g) in 74% yield as an oil. This material was
20 used in the next step without further purification.

¹H NMR (CDCl₃) δ ppm 7.55 (1H, m), 6.77 (1H, d, J=7.4 Hz), 6.61 (1H, d, J=8.1 Hz), 4.66 (2H, d, J=5.3 Hz), 4.38 (2H, q, J=7.1 Hz), 3.46 (1H, t, J=5.2 Hz), 1.41 (3H, t, J=7.1 Hz).

(c) 6-Ethoxy-pyridine-2-carbaldehyde

25 DMSO (0.50 mL, 6.4 mmol) in CH₂Cl₂ (10 mL) was added dropwise to a solution of oxalyl chloride (2M in CH₂Cl₂, 3.1 mL, 6.1 mmol) in CH₂Cl₂ (20 mL) at -60 °C. The resulting mixture was stirred at -60 °C for 10 minutes. (6-Ethoxy-pyridin-2-yl)-methanol (0.85 g, 5.6 mmol) in CH₂Cl₂ (5 mL) and DMSO (4 mL) was added dropwise. The mixture was stirred at -60 °C for 3 h, and was then allowed to warm to -20 °C and Et₃N (6 mL)
30 was added. The resulting solution was stirred at ambient temperature for 40 minutes. Water was added and the mixture was extracted with CH₂Cl₂. The organic phase was washed with brine, dried (Na₂SO₄), and concentrated. Diethyl ether was added to the residue and

insoluble material was removed by filtration. The filtrate was concentrated to yield the title compound (0.60 g) in 70% yield as a solid. This crude product was used in the next step without further purification.

¹H NMR (CDCl₃) δ ppm 9.93 (1H, s), 7.71 (1H, m), 7.53 (1H, d, J=7.1 Hz), 6.94 (1H, d, J=8.3 Hz), 4.46 (2H, d, J=7.1 Hz), 1.42 (3H, t, J=7.1 Hz).

(d) 3-[(6-Ethoxy-pyridin-2-ylmethyl)-amino]-1H-pyrrole-2-carboxylic acid ethyl ester

Acetic acid (0.3 mL) was added to 6-ethoxy-pyridine-2-carbaldehyde (0.59 g, 3.9 mmol) and 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.30 g, 1.9 mmol) in ethanol (10 mL). After 1.5 h, NaCNBH₃ (0.24 g, 3.9 mmol) was added, and the resulting mixture was stirred at ambient temperature for 19 h. The solvent was removed in *vacuo*, ethyl acetate was added to the residue, and insoluble material was removed by filtration. The filtrate was concentrated and the crude product was purified by flash column chromatography (heptane/ethyl acetate gradient; 0 to 35% ethyl acetate), obtaining 0.25 g (45%) of the title product as a solid.

¹H NMR (CDCl₃) δ ppm 8.16 (1H, br s), 7.50 (1H, m), 6.89 (1H, d, J=7.3 Hz), 6.70 (1H, br s), 6.57 (1H, d, J=8.1 Hz), 4.45 (2H, q, J=7.0 Hz), 5.71 (1H, m), 4.38 (5H, m), 1.37 (6H, m);

MS (ESI) m/z 290 (M+1).

(e) 1-(6-Ethoxy-pyridin-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

Ethoxycarbonyl isothiocyanate (0.12 g, 0.90 mmol) was added to 3-[(6-ethoxy-pyridin-2-ylmethyl)-amino]-1H-pyrrole-2-carboxylic acid ethyl ester (0.24 g, 0.82 mmol) in CH₂Cl₂ (5 mL) and the solution was stirred at ambient temperature for 35 minutes. The solvent was evaporated and 0.4M NaOEt in ethanol (3 mL, 1.2 mmol) was added to the residue and the mixture was refluxed for 1 h. More NaOEt (0.4M in ethanol, 1.5 mL, 0.6 mmol) was added and the solution was refluxed for another 1.5 h. The solvent was evaporated, the residue was dissolved in water and the pH adjusted to neutral pH with 1M HCl. The resulting solid was collected, washed, and dried to give crude product. This material was purified by preparative HPLC to yield the title compound (38 mg, 15%) as a solid.

¹H NMR (DMSO-d₆) δ ppm 12.32 (2H, br s), 7.62 (1H, m), 7.29 (1H, d, J=3.0 Hz), 6.78 (1H, d, J=7.3 Hz), 6.64 (1H, d, J=8.3 Hz), 6.13 (1H, d, J=2.8 Hz), 5.65 (2H, s), 4.17 (2H, q, J=7.1 Hz), 1.19 (3H, t, J=7.0 Hz);

^{13}C NMR (DMSO- d_6) δ 173.5, 162.7, 152.8, 152.5, 139.6, 137.2, 127.8, 113.9, 113.6, 109.0, 97.0, 61.0, 53.8, 14.3;
MS (ESI) m/z 303 ($M+1$).

5 **Example 7**

1-Piperidin-3-ylmethyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 3-Formyl-piperidine-1-carboxylic acid tert-butyl ester

DMSO (0.18 mL, 2.6 mmol) in CH_2Cl_2 (5 mL) was added dropwise to a solution of oxalyl chloride (2M in CH_2Cl_2 , 0.65 mL, 1.3 mmol) in CH_2Cl_2 (4 mL) at -78°C . The resulting
10 mixture was stirred at -68°C for 15 minutes. 3-Hydroxymethyl-piperidine-1-carboxylic acid tert-butyl ester (Dean A. Wacker et al. *Bioorganic & Medicinal Chemistry Letters* 2002, 12, 1785-1789) (0.22 g, 1.0 mmol) in CH_2Cl_2 (4 mL) was added dropwise and after 15 min stirring at -78°C , Et_3N (6 mL) was added. The resulting solution was stirred at ambient temperature for 16 h. Water was added and the mixture was extracted with diethyl
15 ether, the organic layer was dried (Na_2SO_4) and concentrated to give the product as an yellow oil (0.20 g, 92% yield). This crude product was used in the next step without further purification.

MS (ESI) m/z 214 ($M+1$).

(b) 3-[(2-Ethoxycarbonyl-1H-pyrrol-3-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester

20 The title compound was obtained as an oil in 30% yield starting from 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.14 g, 0.92 mmol) and 3-formyl-piperidine-1-carboxylic acid tert-butyl ester (0.20 g, 0.92 mmol).

^1H NMR (CDCl_3) δ ppm 8.29 (1H, br s), 6.70 (1H, s), 5.67 (1H, m), 4.27 (2H, m), 3.93
25 (1H, br s), 3.85 (1H, d, $J=13.2$ Hz), 3.10-2.94 (2H, m), 2.83 (1H, m), 2.65 (1H, br s), 1.85 (1H, m), 1.76 (1H, m), 1.64 (1H, m), 1.42 (9H, s), 1.31 (3H, t, $J=6.8$ Hz), 1.22 (1H, m);
MS (ESI) m/z 352 ($M+1$).

(c) 1-Piperidin-3-ylmethyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound was obtained as a solid in 20% yield using 3-[(2-ethoxycarbonyl-1H-pyrrol-3-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester (97 mg, 0.27
30 mmol) and ethoxycarbonyl isothiocyanate (36 mg, 0.27 mmol) using the general procedure B, with the following modifications. After the base mediated cyclization reaction, 6M HCl

(0.3 mL) was added to the reaction followed by heating in a microwave reactor for 4 minutes at 100 °C. The solvent was removed *in vacuo* and the residual solid was purified by preparative HPLC using Atlantis C18 19 x 100 mm, 5 µm column. Gradient with acetonitrile/0.1M ammonium acetate in 5% acetonitrile in MilliQ Water, run from 0% to 50% acetonitrile, in 15 min. Flow rate: 15 ml/min.

¹H NMR (Methanol-d₄) δ ppm 7.23 (1H, d, J=3.2 Hz), 6.21 (1H, d, J=3.2 Hz), 4.46 (1H, m), 4.23 (1H, m), 3.22 (2H, m), 2.86 (2H, m), 2.58 (1H, m), 1.83 (2H, m), 1.62 (1H, m), 1.42 (1H, m);

¹³C NMR (Methanol-d₄) δ 178.8, 154.7, 139.3, 129.5, 115.3, 97.8, 53.7, 48.0, 45.2, 34.5, 27.6, 23.2;

MS (ESI) m/z 265 (M+1).

Example 8

1-Butyl-4-thioxo-1,3,4,5-tetrahydro-2H-pyrrolo[3,2-d]pyrimidin-2-one

Ethoxycarbonyl isothiocyanate (0.13 ml, 1.1 mmol) was added to 3-(butylamino)-1H-pyrrole-2-carboxylic acid ethyl ester (0.23 g, 1.1 mmol) in toluene (5 mL) and the mixture was heated at 90 °C for 1 h. The precipitate was filtered off and washed with hexane. The intermediate product was treated with potassium hydroxide (0.55 g, 9.9 mmol) in water (9 mL), and heated to reflux for 15 h. After cooling to ambient temperature, the pH was adjusted to pH 5 with 12 M HCl. The resultant precipitate was collected by filtration and washed with water. The crude product was purified using preparative HPLC to give the title compound (16 mg, 6%) as a solid.

¹H NMR (DMSO-d₆) δ ppm 12.04 (1 H, br s), 11.95 (1 H, br s), 7.40 (1 H, s), 6.23 (1 H, d, J=2.7 Hz), 3.84 (2 H, t, J=7.2 Hz), 1.66 – 1.56 (2 H, m), 1.32 (2 H, m), 0.89 (3 H, t, J=7.3 Hz);

MS (ESI) m/z 224 (M +1).

Example 9

1-(2-Isopropoxyethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 3-[(2-Isopropoxyethyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester

Trichlorocyanuric acid (1.84 g, 7.93 mmol) was added to a solution of 2-isopropoxyethanol (0.75 g, 7.21 mmol) in CH₂Cl₂ (3 mL). The reaction mixture was

cooled to 0 °C and TEMPO (0.022 g, 0.14 mmol) was carefully added in small portions. The mixture was stirred at r.t. for 20 minutes then filtered through Celite and washed with CH₂Cl₂. The filtrate was kept cold, 0 °C, during filtration. The aldehyde solution was added to a stirred mixture of 3-amino-1H-pyrrole-2-carboxylic acid ester (0.83 g, 5.41 mmol) and HOAc (0.62 mL, 10.8 mmol) at 0 °C in methanol (5 mL). The mixture was stirred for 20 minutes, then NaCNBH₃ (0.34 g, 5.41 mmol) was added. After stirring at r.t for 2 h, the solution was evaporated onto silica and purified by flash column chromatography (heptane/ethyl acetate gradient; 0 to 100% ethyl acetate) to yield the title compound (0.75 g, 58%) as an oil.

¹H NMR (DMSO-d₆) δ ppm 10.72 (1H, br s), 6.76-6.74 (1H, m), 5.66-5.65 (1H, m), 5.34 (1H, br s), 4.17 (2H, q, J=7.0 Hz), 3.59-3.49 (3H, m), 3.15 (2H, q, J=5.6 Hz), 1.26 (3H, t, J=7.0 Hz), 1.10 (3H, s), 1.08 (3H, s);
MS (ESI) m/z 241 (M +1).

(b) 1-(2-Isopropoxyethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound (0.17 g, 23%) was prepared in accordance with the general method B using 3-[(2-isopropoxyethyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester (0.7 g, 2.91 mmol) and ethoxycarbonyl isothiocyanate (0.40 mL, 3.50 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.74 (2H, br s), 7.35 (1H, d, J=2.8 Hz), 6.29 (1H, d, J=3.0 Hz), 4.49 (2H, t, J=6.3 Hz), 3.72 (2H, t, J=6.3 Hz), 3.60-3.58 (1H, m), 1.02 (3H, s), 1.01 (3H, s);

MS (ESI) m/z 254 (M +1).

Example 10

1-(2-Methoxy-2-methylpropyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 3-[(2-Methoxy-2-methylpropyl)amino]-1H-pyrrole-2-carboxylic acid ester

The title compound was obtained as an oil in 75% yield starting from 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.250 g, 1.62 mmol) and 2-methoxy-2-methylpropanal

(US 3,652,579) (0.331 g, 3.24 mmol) using the general procedure A but with the following modifications. After 6 h more 2-methoxy-2-methylpropanal (0.165 g, 1.62 mmol) was added, and the reaction mixture was then stirred overnight.

¹H NMR (DMSO-d₆) δ ppm 10.69 (1H, br s), 6.74 (1H, t, J=3.0 Hz), 5.64 (1H, t, J=2.6 Hz), 5.33 (1H, br s), 4.17 (2H, q, J=7.1 Hz), 3.11 (3H, s), 3.03 (2H, d, J=5.8 Hz), 1.26 (3H, t, J=7.1 Hz), 1.13 (6H, s);
MS (ESI) m/z 241 (M +1).

(b) 1-(2-Methoxy-2-methylpropyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound was obtained as a solid in 3% yield starting from 3-[(2-ethoxy-2-methylpropyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester (0.283 g, 1.18 mmol) and ethoxycarbonyl isothiocyanate (0.13 mL, 1.18 mmol) using the general procedure B but with the following modification. The reaction was run in a microwave reactor for a total of 35 minutes.

¹H NMR (DMSO-d₆) δ ppm 12.29 (1H, br s), 12.17 (1H, br s), 7.30 (1H, d, J=2.76), 6.29 (1H, d, J=2.76), 4.58 (2H, br s), 3.12 (3H, s), 1.21 (6H, s);
MS (ESI) m/z 254 (M +1).

Example 11

1-(2-Ethoxy-2-methylpropyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 2-Bromo-1,1-diethoxy-2-methylpropane

The product was synthesized according to a modified procedure described in US 3,652,579. Bromine water (2.95 mL, 57.6 mmol) was added dropwise to isobutyraldehyde (4.82 g, 66.8 mmol) in ethanol (22 mL) and the resulting mixture was stirred at r.t. for 40 minutes. More bromine water (0.3 mL, 5.86 mmol) was added. The reaction mixture was neutralized by addition of calcium carbonate (3.5 g, 25.3 mmol). The remaining calcium carbonate was filtered off and the filtrate was poured onto an ice-water mixture. The aqueous phase was extracted with CH₂Cl₂, dried (Na₂SO₄), filtered and concentrated. After vacuum distillation, the title product (10.10 g, 67%) was obtained.

¹H NMR ((DMSO-d₆) δ ppm 4.43 (1H, s), 3.80-3.73 (2H, m), 1.64 (6H, s), 1.15 (6H, t, J=7.1 Hz).

(b) 2-Ethoxy-2-methylpropanal

The product was synthesized according to a procedure described in US 3,652,579.

2-Ethoxy-2-methylpropanal (5.63 g, 25 mmol) was added dropwise to potassium bitartrate (2.35 g, 12.5 mmol) in refluxing deionized water (22.5 mL) over 50 minutes. The resulting mixture was refluxed for 70 minutes. The solvent and product were distilled off.

Ammonium sulfate (total 8.5 g) was added to the product-solvent mixture. The mixture was stirred and then the two phases were separated and the upper phase was distilled from calcium chloride obtaining the title product (1.60 g, 55%).

MS (CI) m/z 117 (M +1).

(c) 3-[(2-Ethoxy-2-methylpropyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester

The title compound was obtained as an oil in 63% yield starting from 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.200 g, 1.30 mmol) and 2-ethoxy-2-methyl propionaldehyde (0.292 g, 2.86 mmol) using the general procedure A but with the following modification.

The reaction mixture was stirred at r.t. for 48 h.

¹H NMR (CDCl₃) δ ppm 6.74 (1H, br s), 5.70 (1H, br s), 4.32 (2H, q, J=7.4 Hz), 3.54-3.47 (2H, m), 3.44 (2H, q, J=7.6 Hz), 3.12 (2H, d, J=4 Hz), 1.25 (6H, s), 1.20 (3H, t, J=7.4 Hz), 1.19 (3H, t, J=7.6 Hz);

MS (ESI) m/z 255 (M +1).

(d) 1-(2-Ethoxy-2-methylpropyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

3-(2-Methoxy-2-methyl)-propylamino-1H-pyrrole-2 carboxylic acid ethyl ester (0.200 g, 0.79 mmol) was dissolved in CH₂Cl₂ (2 mL) at r.t. under a nitrogen atmosphere.

Ethoxycarbonyl isothiocyanate (0.12 mL, 1.02 mmol) was added dropwise and the reaction mixture was stirred at r.t. overnight. The solvent was evaporated and sodium ethoxide (1M in ethanol, 0.94 mL, 0.94 mmol) was added and the reaction was heated to 40 °C for 48 h. Water (2 mL) was added and the pH was adjusted to neutral pH with 2M HCl. The precipitate was collected by filtration and was purified by preparative HPLC to give the title compound in 6% yield (0.12 g).

¹H NMR (DMSO-d₆) δ ppm 12.22 (1H, br s), 7.30 (1H, d, J=2.8 Hz), 6.35 (1H, d, J=3 Hz), 4.60 (2H, br s), 3.40-3.34 (3H, m), 1.22 (6H, s), 1.04 (3H, t, J=7.0 Hz);
MS (ESI) m/z 267 (M +1).

5 **Example 12**

1-(Piperidin-4-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one
(a) 4-[(2-(Ethoxycarbonyl)-1H-pyrrol-3-ylamino)-methyl]piperidine-1-carboxylic acid
tert-butyl ester

The title compound (0.156 g, 10%) was prepared in accordance with general method A
10 starting from 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.68 g, 4.4 mmol) and 4-formylpiperidine-1-carboxylic acid tert-butyl ester (P. C. Ting et al., *Bioorganic & Medicinal Chemistry Letters*, 2001, 11, 491-494) (0.98 g, 4.6 mmol).

¹H NMR (DMSO-d₆) δ ppm 10.70 (1H, br s), 6.74 (1H, br s), 5.65 (1H, br s), 4.19 (2H, q, J=7.2 Hz), 3.95 (2H, d, J=12.0 Hz), 2.97 (2H, t, J=6.0 Hz), 2.65 (2H, br s), 1.66 (2H, d, J=12.0 Hz), 1.39 (9H, s), 1.26 (3H, t, J=7.2 Hz), 1.07-0.95 (2H, m);
15

MS (ESI) m/z 352 (M +1).

(b) 1-(Piperidin-4-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

Ethoxycarbonyl isothiocyanate (0.058 g, 0.44 mmol) was added to a stirred solution of 4-[(2-(ethoxycarbonyl)-1H-pyrrol-3-ylamino)-methyl]piperidine-1-carboxylic acid tert-butyl ester (0.156 g, 0.44 mmol) in CH₂Cl₂ (2 mL) and the mixture was stirred at r.t. for 1
20 h. The solvent was removed *in vacuo* and the residue was taken up in ethanol (1 mL) containing sodium (0.015 g, 0.66 mmol). The resulting mixture was heated in a microwave reactor at 120 °C for 10 minutes. 6M HCl (0.5 mL) was added and the reaction mixture was heated again in the microwave at 100 °C for 3 minutes. The pH was adjusted to neutral
25 pH with 2M HCl and the solution was concentrated *in vacuo*. The crude product was purified by preparative HPLC to give the title compound (0.038 g, 14%) as a white solid.
¹H NMR (DMSO-d₆) δ ppm 7.36 (1H, d, J=2.8 Hz), 6.33 (1H, d, J=2.8 Hz), 4.27 (2H, br s), 2.95 (2H, d, J=12.0 Hz), 2.40 (2H, t, J=10.4 Hz), 2.25-2.15 (1H, m), 1.50 (2H, d, J=10.8 Hz), 1.37-1.20 (2H, m);

^{13}C NMR (DMSO- d_6) δ ppm 173.2, 152.9, 137.6, 128.1, 114.0, 97.5, 55.2, 45.5, 34.7, 30.0;

MS (ESI) m/z 265 ($M+1$).

5 **Example 13**

1-[(1-Methylpiperidin-3-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

1-Piperidin-3-ylmethyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one (Example 7) (0.092 g, 0.35 mmol) was dissolved in methanol (2 mL) and formic acid (37%
10 aq., 0.059 mL, 0.7 mmol) was added. After 5 minutes of stirring at r.t. a precipitate had formed. NaCNBH₃ (0.026 g, 0.42 mmol) was added and the mixture was stirred at r.t. for 1 h. The solvent was removed *in vacuo* and the residual solid was purified by preparative HPLC, obtaining the title compound (0.022 g, 22%) as a white solid.

^1H NMR (DMSO- d_6) δ ppm 12.22 (1H, br s), 7.36 (1H, d, $J=2.8$ Hz), 6.33 (1H, s), 4.27
15 (2H, br s), 2.61-2.5 (1H, m), 2.36-2.30 (1H, m), 2.09 (3H, s), 1.93-1.82 (3H, m), 1.65-1.52 (2H, m), 1.44-1.32 (1H, m), 1.16-1.07 (1H, m);

^{13}C NMR (DMSO- d_6) δ ppm 173.2, 152.9, 137.6, 128.1, 114.0, 97.3, 59.2, 56.0, 53.3, 46.7, 34.9, 27.7, 24.7;

MS (ESI) m/z 279 ($M+1$).

20

Example 14

1-[2-Hydroxy-2-(4-methoxyphenyl)ethyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) Methyl {[*tert*-butyl(dimethyl)silyl]oxy}(4-methoxyphenyl)acetate

25 TBDMSCl (1.5 g, 9.94 mmol) and imidazole (1.0 g, 14.6 mmol) were added to a solution of methyl hydroxy-(4-methoxyphenyl)acetate (Teodozyj Kolasa et al., *J. Org. Chem.*, 1987, 22, 4978-4984) (1.3 g, 6.62 mmol) in DMF (8 mL) and the mixture was stirred at r.t. for 2 h. Water was added and the mixture was extracted with diethyl ether. The organic layer was washed with brine, dried (MgSO₄), filtered and concentrated to give the title
30 compound (2.0 g, 97%).

¹H NMR (CDCl₃) δ ppm 7.39 (2H, d, J=8.8 Hz), 6.88 (2H, d, J=8.8 Hz), 5.19 (1H, s), 3.81 (3H, s), 3.69 (3H, s), 0.92 (9H, s), 0.11 (3H, s), 0.03 (3H, s).

(b) *{[tert-Butyl(dimethyl)silyl]oxy}(4-methoxyphenyl)acetaldehyde*

Methyl {[tert-butyl(dimethyl)silyl]oxy}(4-methoxyphenyl)acetate (0.5 g, 1.61 mmol) was dissolved in toluene (10 mL) and cooled to -78 °C under a nitrogen atmosphere. DIBAL (1.0M in toluene, 1.9 mL, 1.93 mmol) was added slowly and the mixture was stirred at -78 °C for 1 h. The reaction mixture was poured onto a mixture of ice (20 g) and CHCl₃ (20 mL). The mixture was stirred at r.t. for 30 minutes. The layers were separated and the water phase extracted with CHCl₃. The organic phase was washed with brine, dried (MgSO₄), filtered and concentrated, obtaining 99% (0.45 g) of the title compound. The product was used directly in the next step without further purification.

¹H NMR (CDCl₃) δ ppm 7.40-7.38 (2H, m), 6.89-6.87 (2H, m), 5.19 (1H, s), 3.81 (3H, s), 0.92 (9H, s), 0.11 (3H, s), 0.03 (3H, s).

(c) *3-{[2-{[tert-Butyl(dimethyl)silyl]oxy}-2-(4-methoxyphenyl)ethyl]amino}-1H-pyrrole-2-carboxylic acid ethyl ester*

The title compound (0.13 g, 19%) was prepared in accordance with general method A using 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.16 g, 1.07 mmol) and {[tert-butyl(dimethyl)silyl]oxy}(4-methoxyphenyl)acetaldehyde (0.3 g, 1.07 mmol). MS (ESI) m/z 417 (M -1).

(d) *1-[2-{[tert-Butyl(dimethyl)silyl]oxy}-2-(4-methoxyphenyl)ethyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one*

The title compound (0.07 g, 90%) was prepared in accordance with general method B using 3-{[2-{[tert-butyl(dimethyl)silyl]oxy}-2-(4-methoxyphenyl)ethyl]amino}-1H-pyrrole-2-carboxylic acid ethyl ester (0.13 g, 0.31 mmol) and ethoxycarbonyl isothiocyanate (0.042 mL, 0.37 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.28-12.23 (2H, m), 7.42 (2H, d, J=8.6 Hz), 7.34-7.33 (1H, m), 6.96 (2H, d, J=8.6 Hz), 6.33 (1H, br s), 5.53-5.50 (1H, m), 4.57 (1H, br s), 4.15 (1H, br s), 3.76 (3H, s), 0.61 (9H, s), 0.31 (3H, s), 0.39 (3H, s); MS (ESI) m/z 432 (M +1).

(e) *1-[2-Hydroxy-2-(4-methoxyphenyl)ethyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one*

Tetra-n-butylammonium fluoride (1M in THF, 1.27 mL, 1.27 mmol) was added to

1-[2-{{*tert*-butyl(dimethyl)silyl}oxy}-2-(4-methoxyphenyl)ethyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-*d*]pyrimidin-4-one (0.065 g, 0.152 mmol) in THF (8 mL). The mixture was stirred at 50 °C overnight. Ethyl acetate was added, and the organic phase was washed with water and brine, dried (MgSO₄), filtered and concentrated. This crude material was purified by preparative HPLC to yield the title compound (0.018 g, 37%) as a solid.

¹H NMR (DMSO-*d*₆) δ ppm 12.20 (2H, br s), 7.40 (2H, d, *J*=8.5 Hz), 7.29 (1H, d, *J*=2.8 Hz), 6.91 (2H, d, *J*=8.8 Hz), 6.27 (1H, d, *J*=2.8 Hz), 5.42-5.41 (1H, m), 5.28-5.24 (1H, m), 4.62-4.58 (1H, m), 4.19-4.16 (1H, m), 3.74 (3H, s);

MS (ESI) *m/z* 316 (*M* -1).

Example 15

1-(2-Methoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-*d*]pyrimidin-4-one

(a) 3-[(2-Methoxybenzyl)amino]-1*H*-pyrrole-2-carboxylic acid ethyl ester

The title compound was obtained as a white solid in quantitative yield starting from 3-amino-1*H*-pyrrole-2-carboxylic acid ethyl ester (0.350 g, 2.27 mmol) and *ortho*-anisaldehyde (0.37 g, 2.71 mmol) using general procedure A but with the following modifications. After stirring overnight, the reaction mixture was evaporated. The crude solid was taken up in CHCl₃, filtered and the solvent was evaporated *in vacuo*, and this crude product was used in the next step without further purification.

MS (ESI) *m/z* 275 (*M* +1).

(b) 1-(2-Methoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-*d*]pyrimidin-4-one

The title compound was obtained as a solid in 16% yield starting from 3-[(2-methoxybenzyl)amino]-1*H*-pyrrole-2-carboxylic acid ethyl ester (0.622 g, 2.27 mmol) and ethoxycarbonyl isothiocyanate (0.26 mL, 2.27 mmol) using general procedure B but with the following modification. The intermediate crude product was dissolved in 1*M* NaOEt (2.27 mL, 2.27 mmol) and was stirred at 80 °C for 3 h.

¹H NMR (DMSO-*d*₆) δ ppm 12.36 (2H, br s), 7.27 (1H, d, *J*=2.8 Hz), 7.27-7.21 (1H, m), 7.06 (1H, d, *J*=8.1 Hz), 6.82 (1H, t, *J*=7.3 Hz), 6.79-6.75 (1H, m), 5.96 (1H, d, *J*=2.8 Hz), 5.61 (2H, s), 3.89 (3H, s);

MS (ESI) *m/z* 288 (*M* +1).

Example 16**1-(3-Methoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one***(a) 3-[(3-Methoxybenzyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester*

5 The title compound was obtained as an oil in 57% (0.508 g) yield and was prepared in accordance with general method A using 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.50 g, 3.24 mmol) and m-anisaldehyde (0.47 mL, 3.89 mmol).

¹H NMR (DMSO-d₆) δ ppm 10.73 (1H, br s), 7.24-7.20 (1H, m), 6.90-6.86 (1H, m), 6.80-6.77 (1H, m), 6.71-6.69 (1H, m), 5.75 (1H, br s), 5.59-5.58 (1H, m), 4.46 (1H, d, J=5.8 Hz), 4.25 (2H, d, J=6.3 Hz), 4.19 (2H, q, J=7.1 Hz), 3.72 (3H, s), 1.26 (3H, t, J=7.1 Hz);
10 MS (ESI) m/z 275 (M +1).

(b) 1-(3-Methoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound was obtained as a solid in 3% (0.014 g) yield and was prepared in accordance with general method B using 3-[(3-methoxybenzyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester (0.494 g, 1.80 mmol) and ethoxycarbonyl isothiocyanate (0.20 mL, 1.18 mmol).
15

¹H NMR (DMSO-d₆) δ ppm 12.41-12.34 (2H, m), 7.29 (1H, d, J=2.7 Hz), 7.23 (1H, t, J=8.0 Hz), 6.93-6.91 (1H, m), 6.86 (1H, d, J=7.8 Hz), 6.83 (1H, dd, J=8.2, 2.4 Hz), 6.14 (1H, d, J=2.8 Hz), 5.67 (2H, s), 3.71 (3H, s);

20 MS (ESI) m/z 288 (M +1).

Example 17**1-(2,4-Dimethoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one***(a) 3-[(2,4-Dimethoxybenzyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester*

25 The title compound was obtained as an oil in 85% (0.838 g) yield and was prepared in accordance with general method A using 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.50 g, 3.24 mmol) and 2,4-dimethoxybenzaldehyde (0.647 g, 3.89 mmol).

¹H NMR (DMSO-d₆) δ ppm 10.69 (1H, br s), 7.14 (1H, d, J=8.3 Hz), 6.71 (1H, t, J=3.0 Hz), 6.54 (1H, d, J=2.3 Hz), 6.44 (1H, dd, J=8.3 Hz), 5.66 (1H, t, J=2.5 Hz), 5.59 (1H, br s), 4.20-4.13 (4H, m), 3.80 (3H, s), 3.73 (3H, s), 1.25 (3H, t, J=7.1 Hz);
30

¹³C NMR (DMSO-d₆) δ ppm 160.9, 159.6, 158.0, 129.2, 124.0, 120.0, 104.2, 98.3, 95.5, 58.3, 55.4, 55.1, 43.6, 14.7;

MS (ESI) m/z 303 (M -1).

(b) *1-(2,4-Dimethoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one*

5 The title compound was obtained as a solid in 14% (0.118 g) yield and was prepared in accordance with general method B using 3-[(2,4-dimethoxybenzyl)amino]-1H-pyrrole-2-carboxylic acid ethyl ester (0.828 g, 2.72 mmol) and ethoxycarbonyl isothiocyanate (0.31 mL, 2.72 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.41 (1H, br s), 12.27 (1H, s), 7.27 (1H, t, J=2.9 Hz), 6.77
10 (1H, d, J=8.3 Hz), 6.61 (1H, d, J=2.3 Hz), 6.41 (1H, dd, J=8.5, 2.4 Hz), 5.95 (1H, t, J=2.3 Hz), 5.54 (2H, s), 3.88 (3H, s), 3.72 (3H, s);

MS (ESI) m/z 318 (M +1).

Example 18

15 **1-[(3-Chloropyridin-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one**

(a) *3-{[(3-chloropyridin-2-yl)methyl]amino}-1H-pyrrole-2-carboxylic acid ethyl ester*

The title compound was obtained as a solid in 91% (0.225 g) yield and was prepared in accordance with general method A using 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester
20 (0.231 g, 1.50 mmol) and 3-chloropyridine-2-carbaldehyde (Nadeem Iqbal et al., *J. Med. Chem.* 1998, 41, 1827-1837) (0.212 g, 1.50 mmol).

¹H NMR (DMSO-d₆) δ ppm 10.81 (1H, br s), 8.53-8.51 (1H, m), 7.94-7.92 (1H, m), 7.40-7.36 (1H, m), 6.77-6.76 (1H, m), 5.74-5.73 (1H, m), 4.43 (1H, d, J=5.5 Hz), 4.20-4.15 (2H, m), 1.30-1.27 (3H, m);

25 MS (ESI) m/z 280 (M +1).

(b) *1-[(3-Chloropyridin-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one*

The title compound was obtained as a solid in 5% (0.011 g) yield and was prepared in accordance with general method B using 3-{[(3-chloropyridin-2-yl)methyl]amino}-1H-
30 pyrrole-2-carboxylic acid ethyl ester (0.215 g, 0.77 mmol) and ethoxycarbonyl isothiocyanate (0.09 mL, 0.77 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.26 (1H, br s), 8.32-8.30 (1H, m), 7.96-7.93 (1H, m), 7.34-7.30 (1H, m), 7.28 (1H, d, J=3.0 Hz), 6.16 (1H, d, J=2.8 Hz), 5.80 (2H, s);
MS (ESI) m/z 293 (M +1).

Example 19

1-{{3-(2-Ethoxyethoxy)pyridin-2-yl}methyl}-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 3-(2-Ethoxyethoxy)-2-methylpyridine

Potassium carbonate (2.20 g, 15.9 mmol) was added to a stirred solution of 3-hydroxy-2-methylpyridine (1.45 g, 13.3 mmol) and 2-chloroethyl ethyl ether (1.75 mL, 15.9 mmol) in DMF (7 mL) and the mixture was stirred at 70 °C overnight. The reaction was not complete and additional 2-chloroethyl ethyl ether (1 equiv.) and potassium carbonate (1 equiv.) were added and the mixture was stirred at 85 °C for 8 h. Water and ethyl acetate were added and the aqueous layer was extracted with ethyl acetate. The organic layer was dried (MgSO₄), filtered and concentrated. The crude product was purified by flash column chromatography (heptane/ethyl acetate gradient; 0 to 50% ethyl acetate), obtaining 1.80 g (75%) the title compound.

¹H NMR (DMSO-d₆) δ ppm 8.03-7.99 (1H, m), 7.33-7.31 (1H, m), 7.18-7.14 (1H, m), 4.13-4.11 (2H, m), 3.73-3.71 (2H, m), 3.52 (2H, q, J=7.0 Hz), 2.35 (3H, s), 1.12 (3H, t, J=6.9 Hz);

MS (ESI) m/z 182 (M +1).

(b) 3-(2-Ethoxyethoxy)pyridine-2-carbaldehyde

A mixture of 3-(2-ethoxyethoxy)-2-methylpyridine (0.506 g, 2.79 mmol) and selenium dioxide (0.31 g, 2.79 mmol) in 1,4-dioxane (10 mL) was heated at 75 °C overnight. After cooling to r.t., the mixture was filtered and the solids were washed with ethyl acetate. The solvent was removed *in vacuo*. The reaction was not complete and the solid was dissolved in 1,4 dioxane (15 mL) and selenium dioxide (0.31 g, 2.79 mmol) was added. The mixture was heated at 110 °C overnight. Ethyl acetate (10 mL) was added and the mixture was filtered. The black solid was washed with ethyl acetate and the filtrate was evaporated *in vacuo*. The crude product was purified by flash column chromatography (heptane/ethyl acetate gradient; 0 to 100% ethyl acetate), obtaining 0.21 g (39%) of the title compound.

¹H NMR (DMSO-d₆) δ ppm 10.23 (1H, s), 8.35 (1H, d, J=4.3 Hz), 7.77 (1H, d, J=8.6 Hz), 7.66-7.62 (1H, m), 4.29 (2H, m), 3.75 (2H, m), 3.55-3.49 (2H, m), 1.14-1.09 (3H, m).

(c) 3-({[3-(2-Ethoxyethoxy)pyridin-2-yl]methyl}amino)-1H-pyrrole-2-carboxylic acid ethyl ester

5 The title compound (0.17 g, 73%) was prepared in accordance with general method A using 3-(2-ethoxyethoxy)pyridine-2-carbaldehyde (0.21 g, 1.08 mmol) and 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.11 g, 0.717 mmol).

¹H NMR (DMSO-d₆) δ ppm 10.76 (1H, s), 8.13-8.12 (1H, m), 7.46-7.44 (1H, m), 7.30-7.28 (1H, m), 6.76-6.75 (1H, m), 6.29 (1H, br s), 5.71-5.70 (1H, m), 4.32-4.31 (2H, m),
10 4.22-4.17 (4H, m), 3.77-3.74 (2H, m), 3.57-3.51 (2H, m), 1.30 (3H, t, J=7.0 Hz), 1.15-1.12 (3H, m);

MS (ESI) m/z 334 (M +1).

(d) 1-({[3-(2-Ethoxyethoxy)pyridin-2-yl]methyl}-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

15 The title compound (0.051 g, 28%) was prepared in accordance with general method B using 3-({[3-(2-ethoxyethoxy)pyridin-2-yl]methyl}amino)-1H-pyrrole-2-carboxylic acid ethyl ester (0.17 g, 0.52 mmol) and ethoxycarbonyl isothiocyanate (0.07 mL, 0.62 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.32-12.19 (2H, m), 7.92 (1H, d, J=4.0 Hz), 7.47 (1H, d, J=7.8 Hz), 7.32-7.14 (2H, m), 5.98 (1H, d, J=2.8 Hz), 5.73 (2H, s), 4.26-4.23 (2H, m),
20 3.78-3.76 (2H, m), 3.55 (2H, q, J=7.1 Hz), 1.14 (3H, t, J=6.9 Hz);

¹³C NMR (DMSO-d₆) δ 173.5, 152.7, 152.4, 143.7, 140.3, 137.8, 127.7, 123.0, 119.0, 113.5, 96.8, 68.2, 65.8, 49.8, 15.1;

MS (ESI) m/z 347 (M +1).

25 Example 20

1-[(6-Oxo-1,6-dihydropyridin-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 3-{{[(6-Oxo-1,6-dihydropyridin-2-yl)methyl]amino}-1H-pyrrole-2-carboxylic acid ethyl ester

30 6-Oxo-1,6-dihydropyridine-2-carbaldehyde (WO 2002/006272) (0.31 g, 2.5 mmol) was dissolved in EtOH (10 mL). 3-Amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.19 g, 1.3 mmol) was added, followed by HOAc (0.14 mL, 2.5 mmol). The mixture was stirred for 75

minutes at r.t. and then NaCNBH₃ (0.16 g, 2.5 mmol) was added. The reaction mixture was stirred at r.t. overnight. The solvent was evaporated *in vacuo* and the crude product was purified by flash column chromatography (CH₂Cl₂/methanol gradient; 0 to 10% methanol), obtaining 0.288 g (85%) of the title product as an oil that crystallized upon standing.

¹H NMR (DMSO-d₆) δ ppm 11.57 (1H, br s), 10.77 (1H, br s), 7.34-7.30 (1H, m), 6.71-6.70 (1H, m), 6.16-6.13 (2H, m), 5.98 (1H, br s), 5.75 (1H, s), 5.64-5.63 (1H, m), 4.20 (2H, q, J=7.1 Hz), 4.09-4.08 (2H, m), 1.27 (3H, t, J=7.1 Hz);

MS (ESI) m/z 262 (M +1).

(b) 1-[(6-Oxo-1,6-dihydropyridin-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

Benzoyl isothiocyanate (0.27 g, 1.6 mmol) dissolved in CH₂Cl₂ (3 mL) was added to 3-{[(6-oxo-1,6-dihydropyridin-2-yl)methyl]amino}-1H-pyrrole-2-carboxylic acid ethyl ester (0.25 g, 0.96 mmol) in CH₂Cl₂ (7 mL). The resulting mixture was stirred at r.t. overnight. The solvent was removed *in vacuo* and the residue was dissolved in methanol (15 mL) and potassium carbonate (0.50 g, 3.6 mmol) was added. The reaction mixture was stirred at 50 °C for 6.5 h. After cooling to r.t., 1M HCl was added dropwise until a neutral pH was obtained. The resulting precipitate was collected, washed with methanol and purified by preparative HPLC to obtain the title compound (0.097 g, 37%) as a solid.

¹H NMR (DMSO-d₆) δ ppm 12.46-12.38 (2H, m), 11.69 (1H, br s), 7.34-7.29 (2H, m), 6.23 (2H, s), 5.75 (1H, br s), 5.49 (2H, s);

¹³C NMR (DMSO-d₆) δ 173.6, 162.7, 152.5, 140.6, 136.7, 128.0, 113.6, 96.6;

MS (ESI) m/z 275 (M +1).

Example 21

1-(1H-Indol-3-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) Ethyl 3-[(1H-indol-3-ylmethyl)amino]-1H-pyrrole-2-carboxylate

A reaction mixture of 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.075 g, 0.49 mmol), indole-3-carboxaldehyde (0.085 g, 0.58 mmol), NaCNBH₃ (0.031 g, 0.49 mmol) and HOAc (0.056 mL, 0.97 mmol) in methanol (3 mL) was stirred at r.t. overnight. The mixture was concentrated *in vacuo* and the crude product-mixture was used in the next step without further purification.

MS (ESI) m/z 284 (M +1).

(b) 1-(1H-Indol-3-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

A crude mixture of ethyl 3-[(1H-indol-3-ylmethyl)amino]-1H-pyrrole-2-carboxylate (max 0.49 mmol) was added to CH₂Cl₂ (5 mL). A few drops of methanol were added to increase solubility. Benzoylisothiocyanate (0.072 g, 0.53 mmol) was added and the mixture was stirred at r.t. for 1 h. The mixture was concentrated *in vacuo*. Ammonia (7N in methanol, 3 mL) was added and the mixture was heated at 80 °C for 2 h. The mixture was concentrated and purified by preparative HPLC, obtaining the title compound (0.030 g, 21%) as a solid.

¹H NMR (DMSO-d₆) δ ppm 12.13 (2H, br s), 11.08 (1H, s), 7.87 (1H, d, J=8.1 Hz), 7.54 - 7.47 (1H, m), 7.38 - 7.30 (1H, m), 7.29 - 7.25 (1H, m), 7.10 - 7.03 (1H, m), 7.01 - 6.93 (1H, m), 6.33 (1H, d, J=2.8 Hz), 5.88 (2H, s);

¹³C NMR (DMSO-d₆) δ ppm 173.14, 152.86, 136.71, 136.42, 128.02, 126.45, 125.83, 121.61, 119.67, 119.16, 114.36, 111.95, 109.51, 97.74, 46.31;

MS (ESI) m/z 295 (M -1).

Example 22

1-(1H-Benzimidazol-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) Ethyl 3-[(1H-benzimidazol-2-ylmethyl)amino]-1H-pyrrole-2-carboxylate

A reaction mixture of 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.77 g, 4.99 mmol), 1H-benzimidazole-2-carboxaldehyde (0.88 g, 5.99 mmol), NaCNBH₃ (0.31 g, 4.99 mmol) and HOAc (0.57 mL, 9.99 mmol) in methanol (15 mL) was stirred at r.t. overnight. The mixture was then heated at 50 °C for 5 h. Cooled to r.t. and evaporated *in vacuo*. The residue was dissolved in ethyl acetate and washed with water. The aqueous phase was extracted with ethyl acetate (twice). The combined organic layers were dried (MgSO₄) and concentrated. The crude product was purified by flash column chromatography (heptane/ethyl acetate (1:0 to 0:1), obtaining 1.15 g (81%) of the title compound.

¹H NMR (DMSO-d₆) δ ppm 12.27 (1H, s), 10.85 (1H, s), 7.63 - 7.37 (2H, m), 7.17 - 7.08 (2H, m), 6.71 (1H, t, J=3.0 Hz), 5.99 (1H, br s), 5.60 (1H, t, J=2.7 Hz), 4.48 (2H, d, J=5.8 Hz), 4.22 (2H, q, J=7.1 Hz), 1.29 (3H, t, J=7.1 Hz);

MS (ESI) m/z 285 ($M + 1$).

(b) *1-(1H-Benzimidazol-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one*

Ethyl 3-[(1H-benzimidazol-2-ylmethyl)amino]-1H-pyrrole-2-carboxylate (0.33 g, 1.16 mmol) was added to CH_2Cl_2 (3 mL) and methanol was added until clear solution was obtained. The solution was stirred at r.t. for 1 h. Benzoyl isothiocyanate (0.73 mL, 0.46 mmol) was added and after stirring at r.t. for 30 minutes the mixture was concentrated. The residue was dissolved in ammonia (7N in methanol, 7 mL) and the mixture was heated at 80 °C in a sealed steel container for 2 h. After cooling to r.t. the precipitated product was filtered and washed through with methanol, diethyl ether and ethyl acetate, obtaining 0.23 g (66%) of the title compound as a white solid.

^1H NMR ($\text{DMSO}-d_6$) δ ppm 12.27 (3H, br s), 7.61 - 7.37 (2H, m), 7.43 - 7.27 (1H, m), 7.18 - 7.07 (2H, m), 6.19 (1H, d, $J=2.8$ Hz), 5.89 (2H, s);

MS (ESI) m/z 298 ($M + 1$).

Example 23

1-[(5-Chloro-1H-indol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

5-Chloro-1H-indole-2-carbaldehyde (0.15 g, 0.76 mmol), NaCNBH_3 (0.040 g, 0.63 mmol) and Et_3N (0.088 mL, 0.63 mmol) was added to a stirred solution of 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester hydrochloride (0.12 g, 0.63 mmol) in methanol (3 mL). The resulting mixture was stirred at r.t. overnight. The reaction mixture was heated to 50 °C. Additional NaCNBH_3 (0.5 equiv.) was added and the mixture was stirred at 50 °C for 3 h. A few drops of HOAc was added, and after 1 h the reaction mixture was cooled to r.t. and was concentrated *in vacuo*. The residue was dissolved in CH_2Cl_2 (2 mL) and methanol (2 mL). Benzoyl isothiocyanate (0.093 mL, 0.69 mmol) was added and after stirring at r.t. for 1 h the mixture was concentrated *in vacuo*. The residue was dissolved in ammonia (7N in methanol, 3 mL) and heated at 80 °C for 2 h. The precipitated product was filtered and washed through with methanol and diethyl ether, followed by purification by preparative HPLC, obtaining 0.063 g (30%) of the title compound as a solid.

¹H NMR (DMSO-d₆) δ ppm 12.41 (2H, br s), 11.51 - 11.04 (1H, m), 7.51 - 7.46 (1H, m), 7.40 - 7.33 (1H, m), 7.33 - 7.30 (1H, m), 7.07 - 6.99 (1H, m), 6.34 - 6.27 (2H, m), 5.87 - 5.80 (2H, m);

¹³C NMR (DMSO-d₆) δ ppm 173.65, 153.05, 136.84, 135.98, 134.74, 129.28, 128.27, 123.97, 121.26, 119.19, 114.24, 113.25, 100.33, 97.17, 47.31;

MS (ESI) m/z 331 (M +1).

Example 24

1-[(5-Fluoro-1H-indol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound was obtained as a solid in 19% (0.038 g) yield and was prepared in accordance with the general method of Example 23 using 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester hydrochloride (0.12 g, 0.63 mmol), 5-fluoro-1H-indole-2-carbaldehyde (0.12 g, 0.76 mmol), NaCNBH₃ (0.040 g, 0.63 mmol, + 0.5 equiv.), Et₃N (0.088 mL, 0.63 mmol) and benzoyl isothiocyanate (0.093 mL, 0.69 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.38 (2H, br s), 11.10 (1H, s), 7.41 - 7.26 (2H, m), 7.25 - 7.11 (1H, m), 6.96 - 6.79 (1H, m), 6.37 - 6.24 (2H, m), 5.83 (2H, s);

¹³C NMR (DMSO-d₆) δ ppm 173.61, 173.81, 158.46, 156.17, 152.94, 136.85, 136.10, 132.94, 128.39, 128.30, 114.19, 112.71, 112.61, 109.54, 109.28, 104.75, 104.51, 100.82, 100.78, 97.23, 47.35;

MS (ESI) m/z 315 (M +1).

Example 25

1-(1H-Indol-6-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound obtained as a solid in 19% (0.035 g) yield and was prepared in accordance with the general method of Example 23 using 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester hydrochloride (0.12 g, 0.63 mmol), 6-formylindole (0.11 g, 0.76 mmol), NaCNBH₃ (0.040 g, 0.63 mmol, + 0.5 equiv.), Et₃N (0.088 mL, 0.63 mmol) and benzoyl isothiocyanate (0.093 mL, 0.69 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.37 (2H, br s), 11.01 (1H, s), 7.53 - 7.43 (1H, m), 7.33 (1H, s), 7.32 - 7.25 (2H, m), 7.09 - 7.03 (1H, m), 6.41 - 6.34 (1H, m), 6.17 (1H, d, J=2.8 Hz), 5.79 (2H, s);

¹³C NMR (DMSO-d₆) δ ppm 173.76, 152.91, 137.06, 136.21, 128.88, 128.27, 127.31, 125.99, 120.35, 118.91, 114.16, 110.24, 101.28, 97.63, 53.32;
MS (ESI) m/z 297 (M +1).

5 **Example 26**

1-(1H-Indol-5-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

The title compound was obtained as a solid in 39% (0.073 g) yield and was prepared in accordance with the general method of Example 23 using 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester hydrochloride (0.12 g, 0.63 mmol), 5-formylindole (0.11 g, 0.76
10 mmol), NaCNBH₃ (0.040 g, 0.63 mmol, + 0.5 equiv.), Et₃N (0.088 mL, 0.63 mmol) and benzoyl isothiocyanate (0.093 mL, 0.69 mmol).

¹H NMR (DMSO-d₆) δ ppm 12.33 (2H, br s), 11.07 (1H, s), 7.52 (1H, s), 7.36 - 7.29 (2H, m), 7.29 - 7.25 (1H, m), 7.21 - 7.12 (1H, m), 6.39 - 6.34 (1H, m), 6.21 - 6.16 (1H, m), 5.77 (2H, s);

15 ¹³C NMR (DMSO-d₆) δ ppm 173.32, 152.51, 136.63, 135.20, 127.83, 127.50, 126.15, 125.81, 120.54, 118.81, 113.82, 111.40, 101.00, 97.28, 52.99;
MS (ESI) m/z 297 (M +1).

Example 27

20 **1-[(5-Fluoro-1H-indol-3-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one**

3-Amino-1H-pyrrole-2-carboxylic acid ethyl ester hydrochloride (0.10 g, 0.52 mmol) was dissolved in methanol (4 mL) and 5-fluoro-1H-indole-3-carboxaldehyde (0.10 g, 0.63 mmol), NaCNBH₃ (0.033 g, 0.52 mmol) and Et₃N (0.073 g, 0.52 mmol) were added. The
25 resulting mixture was stirred at r.t. overnight. Additional NaCNBH₃ (0.01 g) was added and the mixture was heated at 50 °C for 5 h. The reaction mixture was cooled to r.t. and concentrated *in vacuo*. The crude intermediate was dissolved in CH₂Cl₂ (3 mL) and methanol (1 mL). Benzoyl isothiocyanate (0.078 mL, 0.58 mmol) was added and after stirring at r.t. for 1 h the mixture was concentrated *in vacuo*. The residue was dissolved in
30 ammonia (7N in methanol, 3 mL) and heated at 80 °C for 2 h. The solvent was removed *in vacuo* and after purification by preparative HPLC, the title compound (0.035 g, 21%) was obtained as a solid.

¹H NMR (DMSO-d₆) δ ppm 12.29 (2H, s), 11.38 - 11.08 (1H, m), 7.76 - 7.68 (1H, m), 7.65 - 7.59 (1H, m), 7.37 - 7.30 (1H, m), 7.30 - 7.27 (1H, m), 6.96 - 6.86 (1H, m), 6.37 (1H, d, J=2.8 Hz), 5.84 (2H, s);

¹³C NMR (DMSO-d₆) δ ppm 173.09, 158.27, 155.97, 152.81, 136.59, 133.11, 128.14, 128.09, 126.70, 126.59, 114.37, 113.01, 112.91, 109.95, 109.91, 109.86, 109.68, 104.70, 104.46, 97.74, 46.10;

MS (ESI) m/z 315 (M +1).

Example 28

1-(1H-Imidazol-5-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

3-Amino-1H-pyrrole-2-carboxylic acid ethyl ester hydrochloride (0.10 g, 0.52 mmol) was dissolved in methanol (4 mL) and 4-formylimidazole (0.060 g, 0.63 mmol), NaCNBH₃ (0.033 g, 0.52 mmol) and Et₃N (0.073 g, 0.52 mmol) were added. The resulting mixture was stirred at r.t. overnight. The solvent was removed *in vacuo* and the residue was dissolved in CH₂Cl₂ (3 mL) and methanol (1 mL). Benzoyl isothiocyanate (0.078 mL, 0.58 mmol) was added and after stirring at r.t. for 30 minutes the mixture was concentrated *in vacuo*. The residue was dissolved in ammonia (7N in methanol, 3 mL) and heated at 80 °C for 1 h. The precipitated product was filtered and washed with methanol, followed by diethyl ether. The crude product was purified by preparative HPLC, obtaining 0.017 g (13%) of the title compound as a solid.

OC 710/07

¹H NMR (DMSO-d₆) δ ppm 12.49 - 11.78 (3H, m), 7.53 (1H, s), 7.30 (1H, d, J=2.8 Hz), 7.05 (1H, s), 6.36 (1H, d, J=3.0 Hz), 5.54 (2H, s);

MS (ESI) m/z 335 (M +1).

Example 29

1-(1H-Imidazol-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) Ethyl 3-[(1H-imidazol-2-ylmethyl)amino]-1H-pyrrole-2-carboxylate

A reaction mixture with 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.2 g, 1.30 mmol), 2-imidazolecarboxaldehyde (0.15 g, 1.53 mmol), NaCNBH₃ (0.082 g, 1.30 mmol)

and OHAc (0.15 mL, 2.60 mmol) in methanol (5 mL) was stirred at r.t. for 2 h. The solvent was evaporated *in vacuo* and the residue was dissolved in ethyl acetate. Washed with water and the aqueous layer was extracted twice with ethyl acetate. The combined organic layers were dried (MgSO₄), filtered and concentrated. The crude product was purified by flash column chromatography (CH₂Cl₂/methanol gradient; 0 to 20% methanol), obtaining 0.30 g (99%) of the title compound.

¹H NMR (DMSO-d₆) δ ppm 10.84 (1H, br s), 7.03 (2H, s), 6.79 - 6.64 (1H, m), 5.76 (1H, br s), 5.68 - 5.57 (1H, m), 4.29 (2H, d, J=5.8 Hz), 4.19 (2H, q, J=7.1 Hz), 3.16 (1H, s), 1.26 (3H, t, J=7.1 Hz);

MS (ESI) m/z 235 (M +1).

(b) *1-(1H-Imidazol-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one*

Benzoyl isothiocyanate (0.19 mL, 1.41 mmol) was added to a stirred solution of ethyl 3-[(1H-imidazol-2-ylmethyl)amino]-1H-pyrrole-2-carboxylate (0.3 g, 1.28 mmol) in CH₂Cl₂ (4 mL) and methanol (2 mL) and the mixture was stirred at r.t. for 1h. The solvent was evaporated *in vacuo* and the residue was dissolved in ammonia (7N in methanol, 7 mL) and heated at 80 °C for 1h. The crude product was filtrated and purified by preparative HPLC, obtaining 0.110 g, (35%) of the title compound as a white solid.

¹H NMR (DMSO-d₆) δ ppm 12.46 - 12.14 (2H, m), 11.81 (1H, br s), 7.28 (1H, d, J=2.5 Hz), 6.99 (1H, s), 6.79 (1H, s), 6.13 (1H, d, J=2.8 Hz), 5.67 (2H, s);

MS (ESI) m/z 248 (M +1).

Example 30

1-[(5-Chloro-1H-benzimidazol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) *Ethyl 3-{[(5-chloro-1H-benzimidazol-2-yl)methyl]amino}-1H-pyrrole-2-carboxylate*

A reaction mixture of 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.077 g, 0.5 mmol), NaCNBH₃ (0.057 g, 0.9 mmol) and HOAc (0.030 g, 0.5 mmol) in methanol (4 mL) were stirred at r.t. for 5 minutes before 5-chloro-1H-benzimidazole-2-carbaldehyde (0.144 g, 0.8 mmol) was added, followed by addition of CH₂Cl₂ (1 mL) and DMF (0.4 mL). The resulting mixture was allowed to stir for 16 h at r.t. under nitrogen atmosphere. Additional 5-chloro-1H-benzimidazole-2-carbaldehyde (0.030 g) and NaCNBH₃ (0.015 g) were added

and the reaction mixture was stirred at 4h. The reaction mixture was neutralized with 2M NaOH and diluted with ethyl acetate. Extracted with water and the organic layer was dried (Na_2SO_4) and concentrated. The crude product was purified by flash column chromatography (heptane/ethyl acetate gradient; 0 to 100% ethyl acetate), obtaining 0.074 g (46%) of the title product.

^1H NMR (CDCl_3) δ ppm 8.28 (1H, br s), 7.52 (1H, s), 7.44 (1H, d, $J=2.0$ Hz), 7.22 (1H, d, $J=2.0$ Hz), 6.66 (1H, s), 5.61 (1H, t, $J=2.4$ Hz), 4.67 (2H, s), 4.27 (2H, m), 1.33 (3H, t, $J=6.8$ Hz);

MS (ESI) m/z 319 ($M+1$).

(b) 1-[(5-Chloro-1H-benzimidazol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

A solution of ethyl 3-[(5-chloro-1H-benzimidazol-2-yl)methyl]amino}-1H-pyrrole-2-carboxylate (0.074 g, 0.23 mmol) in CH_2Cl_2 was stirred for 5 minutes. DMF (0.2 mL) was added to enhance solubility. Benzoyl isothiocyanate (0.045 g, 0.28 mmol) was added and the mixture was stirred for 1 h then concentrated *in vacuo*. The crude intermediate was taken up in ammonia (7N in methanol, 2 mL) and was stirred at 70 °C for 1.5 h in a sealed vessel. After cooling to r.t. the precipitated product was collected by vacuum filtration, washed with diethyl ether and dried, obtaining 0.033 g (43%) of the title product as a white solid.

^1H NMR ($\text{DMSO}-d_6$) δ ppm 12.33 (3H, br s), 7.48 (2H, m), 7.32 (1H, s), 7.17 (1H, d, $J=8.0$ Hz), 6.21 (1H, s), 5.88 (2H, s);

MS (ESI) m/z 332 ($M+1$).

Example 31

1-[(4,5-Dimethyl-1H-benzimidazol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) 4,5-Dimethyl-1H-benzimidazole-2-carbaldehyde

3,4-Dimethylbenzene-1,2-diamine (0.409 g, 3.0 mmol) and dichloroacetic acid (0.768 g, 6.0 mmol) in 4N HCl (10 mL) were heated at 100 °C for two days. After cooling to r.t. the mixture was filtrated and the mother liquor was extracted with chloroform (4 times). The pH was set to 12 using 2M NaOH and the resulting white precipitate was collected by filtration. The crude product was used in the next step without further purification.

MS (ESI) m/z 175 ($M + 1$).

(b) *Ethyl 3-{[(4,5-dimethyl-1H-benzimidazol-2-yl)methyl]amino}-1H-pyrrole-2-carboxylate*

3-Amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.077 g, 0.5 mmol), 4,5-dimethyl-1H-benzimidazole-2-carbaldehyde (0.130 g, 0.75 mmol) and HOAc (0.045 g, 0.75 mmol) were stirred in methanol (4 mL) followed by addition of NaCNBH₃. The resulting mixture was stirred at r.t. for 16 h. The reaction mixture was neutralized with 2M NaOH and the solvent was removed *in vacuo*. The residue was dissolved in ethyl acetate and extracted with water. The organic layer was dried (Na₂SO₄) and concentrated. The crude product was purified by flash column chromatography (heptane/ethyl acetate gradient; 0 to 100% ethyl acetate), obtaining a white solid, 0.043 g (27%) of the title compound, as a tautomeric mixture (1:1).

¹H NMR (DMSO-d₆) δ ppm 12.15 (1H, s), 12.04 (1H, s), 10.81 (2H, s), 7.26 (1H, d, $J=8.4$ Hz), 7.12 (1H, d, $J=8.0$ Hz), 6.94 (2H, d, $J=8.4$ Hz), 6.72 (2H, s), 5.93 (2H, s), 5.70 (1H, s), 5.61 (1H, s), 4.44 (4H, dd, $J=9.2, 6.0$ Hz), 4.22 (4H, m), 2.44 (3H, s), 2.37 (3H, s), 2.29 (6H, s), 1.30 (6H, m);

MS (ESI) m/z 313 ($M + 1$).

(c) *1-[(4,5-Dimethyl-1H-benzimidazol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one*

Ethyl 3-{[(4,5-dimethyl-1H-benzimidazol-2-yl)methyl]amino}-1H-pyrrole-2-carboxylate (0.043 g, 0.14 mmol) was dissolved in CH₂Cl₂ (1.5 mL) and benzoyl isothiocyanate (0.026 g, 0.16 mmol) was added. After stirring at r.t. for 1 h the solvent was removed *in vacuo*. The residue was dissolved in ammonia (7N in methanol) and heated to 70 °C for 2 h in a sealed vessel. The solvent was removed *in vacuo* and the residue was purified by preparative HPLC, obtaining a white solid, 0.008 g (18%) of the title compound, as a tautomeric mixture (1:1).

¹H NMR (DMSO-d₆) δ ppm 12.18 (2H, s), 11.98 (2H, s), 7.28 (2H, t, $J=2.4$ Hz), 7.19 (1H, d, $J=8.0$ Hz), 7.08 (1H, d, $J=8.0$ Hz), 6.92 (2H, t, $J=8.0$ Hz), 6.14 (2H, dd, $J=8.0, 2.8$ Hz), 5.89 (2H, s), 5.86 (2H, s), 2.41 (3H, s), 2.37 (3H, s), 2.29 (3H, s), 2.28 (3H, s);

MS (ESI) m/z 326 ($M + 1$).

Exemple 32

7-Bromo-1-isobutyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one*(a) 3-Amino-4-bromo-1H-pyrrole-2-carboxylic acid ethyl ester*

3-Amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.92 g, 0.6 mmol) was dissolved in HOAc (1 mL) and bromine water (0.96 g, 0.6 mmol) was added. The mixture was stirred for 1 h at r.t., and the resulted white precipitate was collected by filtration and washed with diethyl ether. The title compound (0.136 g, 97%) was obtained as a white solid and used in the next step without further purification.

¹H NMR (DMSO-d₆) δ ppm 11.67 (1H, s), 7.05 (1H, s), 6.02 (2H, br s), 4.25 (2H, q, J=7.2 Hz), 1.28 (3H, t, J=7.2 Hz);

MS (ESI) m/z 233 (M +1).

(b) 4-Bromo-3-(isobutylamino)-1H-pyrrole-2-carboxylic acid ethyl ester

3-Amino-4-bromo-1H-pyrrole-2-carboxylic acid ethyl ester (0.136 g, 0.58 mmol) and isobutyraldehyde (0.067 g, 0.93 mmol) were stirred in CH₂Cl₂/MeOH (1:1, 3 mL) at r.t. for 2 h. NaCNBH₃ (0.065 g, 1.04 mmol) and HOAc (0.035 g, 0.58 mmol) were added and the mixture was stirred at r.t. for 2 h and then stirred at 50 °C for 16 h. Additional isobutyraldehyde (1 equiv.) and NaCNBH₃ (0.5 equiv.) were added and the mixture was continued stirring at 50 °C overnight. The reaction mixture was neutralized with 2 M NaOH solution and the solvents were removed *in vacuo*. The residue was taken up in ethyl acetate and extracted with water. The organic layer was dried (Na₂SO₄) and concentrated. The crude product was purified by flash chromatography (heptane/ethyl acetate gradient, 0 to 30% ethyl acetate), obtaining the title compound (0.040 g, 24%) as a white solid.

¹H NMR (CDCl₃) δ ppm 6.75 (1H, s), 4.32 (2H, q, J=7.2 Hz), 3.28 (2H, d, J=6.8 Hz), 1.85 (1H, m), 1.35 (3H, t, J=7.2 Hz), 0.97 (3H, d, J=6.8 Hz);

MS (ESI) m/z 289 (M +1).

(c) 7-Bromo-1-isobutyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

Benzoyl isothiocyanate (0.034 g, 0.2 mmol) was added to a solution of 4-bromo-3-(isobutylamino)-1H-pyrrole-2-carboxylic acid ethyl ester (0.050 g, 0.17 mmol) in CH₂Cl₂ and the resulting mixture was stirred at r.t. for 1 h before it was concentrated *in vacuo*. The residue was dissolved in ammonia (7M in methanol, 1.5 mL) and stirred at 70 °C for 4 h. Additional ammonia (7M in methanol, 1 mL) was added and the mixture was stirred at 80 °C for 5 h. The mixture was cooled to r.t. and the precipitated product was filtered and

washed with diethyl ether. After recrystallization with methanol the title compound (0.028 g, 55%) was obtained as a white solid.

¹H NMR (DMSO-d₆) δ ppm 12.57 (2H, br s), 7.58 (1H, s), 4.92 (1H, br s), 4.42 (1H, br s), 2.39 (1H, m), 0.94 (6H, d, J=6.4 Hz);

5 MS (ESI) m/z 303 (M +1).

Example 33

1-(3-Chlorophenyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one

(a) Ethyl 3-[(3-chlorophenyl)amino]-1H-pyrrole-2-carboxylate

10 A mixture of 3-amino-1H-pyrrole-2-carboxylic acid ethyl ester (0.20 g, 1.3 mmol), 3-bromochlorobenzene (0.30 g, 1.6 mmol), Pd₂(dba)₃ (0.048 g, 0.052 mmol), rac-BINAP (0.048 g, 0.078 mmol) and cesium carbonate (0.59 g, 1.8 mmol) was heated at 100 °C in a sealed microwave vial under nitrogen atmosphere overnight. Additional Pd₂(dba)₃ (0.10 g, 0.11 mmol) and R,S-BINAP (0.11 g, 0.18 mmol) were added and the reaction was
15 continued stirring at 100 °C overnight. Additional 3-bromochlorobenzene (0.15 g), Pd₂(dba)₃ (0.098 g) and R,S-BINAP (0.098 g) were added and the reaction was continued stirring at 100 °C for three more days. The reaction mixture was poured into ethanol and the resulting solution was filtered. The filtrate was evaporated *in vacuo* and the residue was purified by flash column chromatography (heptane/ethyl acetate gradient; 0 to 30% ethyl
20 acetate), obtaining 0.052 g (15%) of the title compound.

¹H NMR (CDCl₃) δ ppm 8.42 (1H, br s), 7.17 (1H, t, J=8.0 Hz), 7.14 (1H, t, J=2.0 Hz), 6.96-6.94 (1H, m), 6.87-6.84 (1H, m), 6.82 (1H, t, J=3.0 Hz), 6.32 (1H, t, J=3.0 Hz), 4.35 (2H, q, J=7.2 Hz), 1.38 (3H, t, J=7.1 Hz);

MS (ESI) m/z 263 (M -1).

25 (b) 1-(3-Chlorophenyl)-2-thioxo-1,2,3,5-tetrahydro--pyrrolo[3,2-d]pyrimidin-4-one

Benzoyl isothiocyanate (0.035 g, 0.22 mmol) in CH₂Cl₂ (0.5 mL) was added to ethyl 3-[(3-chlorophenyl)amino]-1H-pyrrole-2-carboxylate (0.052 g, 0.20 mmol) in CH₂Cl₂ (0.5 mL). The mixture was stirred at r.t. overnight. Additional benzoyl isothiocyanate (0.035 g + 0.035 g + 0.035 g) were added over 6 h and the reaction was continued stirring at 50 °C for
30 3 days. The solvent was removed *in vacuo* and the residue was dissolved in ammonia (7N in methanol, 4 mL). The reaction was heated at 50 °C for 2 h. The crude product was

purified by preparative chromatography, obtaining 0.009 g (15%) of the title compound as a solid.

^1H NMR (DMSO- d_6) δ ppm 12.41 (2H, br s), 7.60-7.57 (3H, m), 7.41-7.37 (1H, m), 7.25 (1H, d, $J=2.8$ Hz), 5.36 (1H, d, $J=2.8$ Hz);

^{13}C NMR (DMSO- d_6) δ ppm 173.8, 152.8, 141.5, 137.9, 133.5, 131.,3 129.0, 128.6, 127.7, 127.3, 113.3, 96.7;

MS (ESI) m/z 278 ($M+1$).

Screens

Methods for the determination of MPO inhibitory activity are disclosed in patent application WO 02/090575. The pharmacological activity of compounds according to the invention was tested in the following screen in which the compounds were either tested alone or in the presence of added tyrosine:

Assay buffer: 20 mM sodium/potassium phosphate buffer pH 6.5 containing 10 mM taurine and 100 mM NaCl.

Developing reagent: 2 mM 3,3',5,5'-tetramethylbenzidine (TMB), 200 μM KI, 200 mM acetate buffer pH 5.4 with 20 % DMF.

To 10 μl of diluted compounds in assay buffer, 40 μl of human MPO (final concentration 2.5 nM), with or without 20 μM tyrosine (final concentration, if present, 8 μM), was added and the mixture was incubated for 10 minutes at ambient temperature. Then 50 μl of H_2O_2 (final concentration 100 μM), or assay buffer alone as a control, were added. After incubation for 10 minutes at ambient temperature, the reaction was stopped by adding 10 μl 0.2 mg/ml of catalase (final concentration 18 $\mu\text{g/ml}$). The reaction mixture was left for an additional 5 minutes before 100 μl of TMB developing reagent was added. The amount of oxidised 3,3',5,5'-tetramethylbenzidine formed was then measured after about 5 minutes


using absorbance spectroscopy at about 650 nM. IC₅₀ values were then determined using standard procedures.

When tested in at least one version of the above screen, the compounds of Examples 1 to 32 gave IC₅₀ values of less than 60 μ M, indicating that they are expected to show useful therapeutic activity. Representative results are shown in the following Table.

Compound	Inhibition of MPO (in the presence of tyrosine) IC₅₀ μM
Example 2	0.26
Example 5	0.22
Example 11	1.1

Claims

1. A compound of formula (I)



wherein:

at least one of X and Y represents S, and the other represents O or S;

L represents a direct bond or C1 to 7 alkylene, said alkylene optionally incorporating a
 10 heteroatom selected from O, S(O)_n and NR⁶, said alkylene optionally incorporating one or
 two carbon-carbon double bonds, and said alkylene being optionally substituted by one or
 more substituents selected independently from OH, halogen, CN and NR⁴R⁵, C1 to 6 alkyl
 and C1 to 6 alkoxy, said alkoxy optionally incorporating a carbonyl adjacent to the
 oxygen;

15 n represents an integer 0, 1 or 2;

R¹ represents hydrogen, or

i) a saturated or partially unsaturated 3 to 7 membered ring optionally incorporating one
 or two heteroatoms selected independently from O, N and S, and optionally incorporating a
 carbonyl group, optionally substituted by one or more substituents independently selected
 20 from halogen, SO₂R⁹, SO₂NR⁹R¹⁰, OH, C1 to 7 alkyl, C1 to 7 alkoxy, CN, CONR²R³,
 NR²COR³ and COR³, said alkoxy being optionally further substituted by C1 to 6 alkoxy
 and said alkoxy optionally incorporating a carbonyl adjacent to the oxygen, and said alkyl
 being optionally further substituted by hydroxy or C1 to 6 alkoxy and said alkyl or alkoxy
 optionally incorporating a carbonyl adjacent to the oxygen or at any position in the alkyl;

25 or

ii) an aromatic ring system selected from phenyl, biphenyl, naphthyl or a monocyclic or bicyclic heteroaromatic ring structure containing 1 to 3 heteroatoms independently selected from O, N and S, said aromatic ring system being optionally substituted by one or more substituents independently selected from halogen, SO_2R^9 , $\text{SO}_2\text{NR}^9\text{R}^{10}$, OH, C1 to 7 alkyl, C1 to 7 alkoxy, CN, CONR^2R^3 , NR^2COR^3 and COR^3 ; said alkoxy being optionally further substituted by C1 to 6 alkoxy and said alkoxy optionally incorporating a carbonyl adjacent to the oxygen, and said alkyl being optionally further substituted by hydroxy or C1 to 6 alkoxy and said alkyl or alkoxy optionally incorporating a carbonyl adjacent to the oxygen or at any position in the alkyl;

R^{12} represents hydrogen or halogen or a carbon optionally substituted with one to three halogen atoms;

at each occurrence, R^2 , R^3 , R^4 , R^5 , R^6 , R^9 and R^{10} independently represent hydrogen, C1 to 6 alkyl or C1 to 6 alkoxy said alkoxy optionally incorporating a carbonyl adjacent to the oxygen, said alkyl being optionally further substituted by halogen, C1 to 6 alkoxy, CHO, C2 to 6 alkanoyl, OH, CONR^7R^8 and NR^7COR^8 ;

or the groups NR^2R^3 , NR^4R^5 and NR^9R^{10} each independently represent a 5 to 7 membered saturated azacyclic ring optionally incorporating one additional heteroatom selected from O, S and NR¹¹, said ring being optionally further substituted by halogen, C1 to 6 alkoxy, CHO, C2 to 6 alkanoyl, OH, CONR^7R^8 and NR^7COR^8 ;

at each occurrence R^7 , R^8 and R^{11} independently represent hydrogen or C1 to 6 alkyl, or the group NR^7R^8 represents a 5- to 7-membered saturated azacyclic ring optionally incorporating one additional heteroatom selected from O, S and NR¹¹;

and pharmaceutically acceptable salts thereof;

with the proviso that the compounds 1- β -D-ribofuranosyl-2-oxopyrrolo[3,2-d]pyrimidine-4(3H,5H)-thione, 1-(2,3,5-tri-O-benzoyl-1- β -D-ribofuranosyl)-2-oxopyrrolo[3,2-d]pyrimidine-4(3H,5H)-thione and 5,7-dimercapto-1,4,6-triazaindene are disclaimed.

2. A compound according to claim 1, wherein R^1 represents hydrogen.

3. A compound according to claim 1 or 2, wherein X represents S and Y represents O.
4. A compound according to claim 1 or 2, wherein Y represents S and X represents O.
- 5 5. A compound according to any one of claims 1 to 4, wherein L is a direct bond or represents C1 to 7 alkylene; said alkylene being optionally substituted by one or more C1 to 6 alkoxy.
6. A compound according to any one of claims 1 to 4, wherein L represents C1 to 3
10 alkylene; said alkylene being optionally substituted by one or more C1 to 6 alkoxy.
7. A compound according to any one of claims 1 to 6, wherein R¹ represents a saturated or partially unsaturated 3 to 7 membered ring optionally incorporating one or two heteroatoms selected independently from O, N and S, and optionally incorporating a
15 carbonyl group; said ring being optionally substituted by one or more substituents independently selected from halogen, C1 to 6 alkyl and C1 to 6 alkoxy, said alkoxy being optionally further substituted by C1 to 6 alkoxy.
8. A compound according to any one of claims 1 to 6, wherein R¹ represents an aromatic
20 ring system selected from phenyl, biphenyl, naphthyl or a five- or six-membered heteroaromatic ring containing 1 to 3 heteroatoms independently selected from O, N and S, said aromatic ring being optionally substituted by one or more substituents independently selected from halogen, C1 to 6 alkyl and C1 to 6 alkoxy, said alkoxy being optionally further substituted by C1 to 6 alkoxy.
- 25 9. A compound according to claim 1, wherein L represents an optionally substituted C1 to 3 alkylene and R¹ represents a saturated or partially unsaturated 3- to 7-membered ring optionally incorporating one or two heteroatoms selected independently from O, N and S, and optionally incorporating a carbonyl group, said ring being optionally substituted by
30 one or more substituents independently selected from halogen, C1 to 6 alkyl and C1 to 6 alkoxy, said alkoxy being optionally further substituted by C1 to 6 alkoxy.

10. A compound according to claim 1, wherein L represents optionally substituted C1 to 3 alkylene and R¹ represents an aromatic ring system selected from phenyl, biphenyl, naphthyl or a five- or six-membered heteroaromatic ring containing 1 to 3 heteroatoms independently selected from O, N and S, said aromatic ring being optionally substituted by
5 one or more substituents independently selected from halogen, C1 to 6 alkyl and C1 to 6 alkoxy, said alkoxy being optionally further substituted by C1 to 6 alkoxy.

11. A compound according to claim 1, wherein X represents S, Y represents O, L represents optionally substituted C1 to 3 alkylene and R¹ represents optionally substituted
10 phenyl.

12. A compound according to claim 1, wherein X represents S, Y represents O, L represents optionally substituted C1 to 3 alkylene and R¹ represents optionally substituted pyridyl.
15

13. A compound according to claim 1, wherein X represents S, Y represents O, L represents C1 to 3 alkylene, substituted with C1 to 6 alkoxy and R¹ represents hydrogen.

14. A compound, said compound being:
20 1-butyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-isobutyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(pyridin-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(2-fluoro-benzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[2-(2-methoxyethoxy)-3-propoxybenzyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-
25 d]pyrimidin-4-one;
1-(6-ethoxy-pyridin-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-piperidin-3-ylmethyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-butyl-4-thioxo-1,3,4,5-tetrahydro-2H-pyrrolo[3,2-d]pyrimidin-2-one;
30 1-(2-isopropoxyethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(2-methoxy-2-methylpropyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

- 1-(2-ethoxy-2-methylpropyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(piperidin-4-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(1-methylpiperidin-3-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
5 1-[2-hydroxy-2-(4-methoxyphenyl)ethyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(2-methoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(3-methoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(2,4-dimethoxybenzyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
10 1-[(3-chloropyridin-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-{[3-(2-ethoxyethoxy)pyridin-2-yl]methyl}-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(6-oxo-1,6-dihydropyridin-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
15 1-(1H-indol-3-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(1H-benzimidazol-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(5-chloro-1H-indol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
20 1-[(5-fluoro-1H-indol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(1H-indol-6-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(1H-indol-5-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
25 1-[(5-fluoro-1H-indol-3-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(1H-imidazol-5-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-(1H-imidazol-2-ylmethyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
1-[(5-chloro-1H-benzimidazol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;
30 1-[(4,5-dimethyl-1H-benzimidazol-2-yl)methyl]-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one;

7-bromo-1-isobutyl-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one; and
1-(3-chlorophenyl)-2-thioxo-1,2,3,5-tetrahydro-pyrrolo[3,2-d]pyrimidin-4-one.

and pharmaceutically acceptable salts thereof.

5 15. A compound of formula (I), according to any one of Claims 1 to 14, or a
pharmaceutically acceptable salt thereof, for use as a medicament.

16. A pharmaceutical composition comprising a compound of formula (I) according to any
one of Claims 1 to 14, or a pharmaceutically acceptable salt thereof, optionally in admixture
10 with a pharmaceutically acceptable adjuvant, diluent or carrier.

17. A method of treating, or reducing the risk of, diseases or conditions in which
inhibition of the enzyme MPO is beneficial which comprises administering to a person
suffering from or at risk of, said disease or condition, a therapeutically effective amount of
15 a compound of formula (I), as defined in any one of Claims 1 to 14, or a pharmaceutically
acceptable salt thereof.

18. A method of treating, or reducing the risk of neuroinflammatory disorders which
comprises administering to a person suffering from or at risk of, said disease or condition,
20 a therapeutically effective amount of a compound of formula (I), as defined in any one of
Claims 1 to 14, or a pharmaceutically acceptable salt thereof.

19. The method according to claim 18, wherein said neuroinflammatory disorder is
multiple sclerosis.

25

20. The method according to claim 17, wherein said disease or condition is
atherosclerosis.

30

21. The method according to claim 17, wherein said disease or condition is COPD.

22. The use of a compound of formula (I) as defined in any one of Claims 1 to 14, or a
pharmaceutically acceptable salt thereof, in the manufacture of a medicament, for the

treatment or prophylaxis of diseases or conditions in which inhibition of the enzyme MPO is beneficial.

23. The use of a compound of formula (I) as defined in any one of Claims 1 to 14, or a
5 pharmaceutically acceptable salt thereof, in the manufacture of a medicament, for the treatment or prophylaxis of neuroinflammatory disorders.


24. The use according to claim 23, wherein said neuroinflammatory disorder is multiple sclerosis.

10

25. The use according to claim 22, wherein said disease or condition is atherosclerosis.

26. The use according to claim 22, wherein said disease or condition is COPD.

15 27. A process for the preparation of a compound of formula (I), as defined in any one of Claims 1 to 14, or a pharmaceutically acceptable salt, enantiomer, diastereomer or racemate thereof, wherein the process comprises reaction of a compound of formula (II)



20

wherein R^1 and L are as defined in Claim 1 and R represents C1 to 6 alkoxy or NH_2 ;
with a C1 to 6 alkoxy carbonyl isothiocyanate or with a phenyl carbonyl isothiocyanate;
and where necessary converting the resultant compound of formula (I), or another salt thereof,
25 into a pharmaceutically acceptable salt thereof; or converting the resultant compound of

formula (I) into a further compound of formula (I); and where desired converting the resultant compound of formula (I) into an optical isomer thereof.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2005/001835

A. CLASSIFICATION OF SUBJECT MATTER

IPC: see extra sheet

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: A61K, C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, CHEM ABS DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 03089430 A1 (ASTRAZENECA AB), 30 October 2003 (30.10.2003), claims 1-11, abstract, RN:618913-14-7, 618913-17-0, 618913-21-6, 618913-22-7, 618913-29-4, 618913-32-9, 618913-16-9, 139460-82-5, 618913-20-5, 618913-24-9, 618913-25-0, 618913-26-1, 618913-27-2,	1-3,5-11, 13-16,22-26
X	618913-28-3, 618913-30-7, 618913-31-8 --	1-3,5-11, 13-16,22-26
X	WO 9940091 A1 (AMGEN INC.), 12 August 1999 (12.08.1999), figure 3	27
A	figure 3, abstract, RN:237435-30-2 --	1-26

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

15 February 2006

Date of mailing of the international search report

21-02-2006

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Eva Johansson/EÖ
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2005/001835

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 0059449 A2 (EURO-CELTIQUE S.A.), 12 October 2000 (12.10.2000), RN:162278-87-7, 162278-88-8, 162278-90-2, 162278-93-5, 162278-94-6, 300783-45-3, 300783-48-6, 300783-53-3 --	1-27
P,A	WO 2005037835 A1 (ASTRAZENECA AB), 28 April 2005 (28.04.2005), claims 1-9, abstract --	1-27
A	WO 03000694 A1 (ALMIRALL PRODESFARMA S.A.), 3 January 2003 (03.01.2003), RN:480992-38-9, 480992-39-0, 480992-40-3, 480992-41-4, 480992-42-5, 480993-71-3 --	1-27
A	WO 03082873 A1 (ALMIRALL PRODESFARMA, S.A.), 9 October 2003 (09.10.2003) --	1-27
A	SUDHAKAR RAO, T. ET AL, "Synthesis of 5,7-Disubstituted-4-Beta-D-ribofuranosylpyrazolo(4,3 -d)-pyrimidines and 2,4-Disubstituted-1-Beta-D-ribofuranosylpyrrolo(3,2- d)-pyrimidines as Congeners of Uridine and Cytidine", J. Heterocyclic Chem., 1992, Volume 29, page 343 - page 354, compound 27 and 28, RN:142545-01-5 --	1-27
A	IMAI, KIN-ICHI ET AL, "Studies on Nucleic Acid Antagonists. VII. Synthesis and Characterization of 1,4,6-Triazaindenenes (5H-Pyrrolo(3,2-d)pyrimidines)", Chem. Pharm. Bull., 1964, Volume 12, no. 9, page 1030 - page 1042, page 1039, compound (PPy-XX), RN:91184-33-7 --	1-27
P,A	WO 2005042534 A2 (CV THERAPEUTICS, INC.), 12 May 2005 (12.05.2005), abstract, claims 1-16,18,26-27, 32 --	1-27

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2005/001835**C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 9618400 A1 (EUROCELTIQUE, S.A.), 20 June 1996 (20.06.1996) -- -----	1-27

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2005/001835

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 17-21
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 17-21 relate to a method of treatment of the human or animal body by surgery or by therapy, as well as diagnostic
.../...
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2005/001835

Box II.1

methods /Rule 39.1(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2005/001835

INTERNATIONAL PATENT CLASSIFICATION (IPC) :

C07D 487/04 (2006.01)

A61K 31/519 (2006.01)

A61P 9/10 (2006.01)

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/SE 2005/001835

WO	03089430	A1	30/10/2003	AT	292371	T	15/04/2005
				AU	2003224548	A	00/00/0000
				BR	0309012	A	01/02/2005
				CA	2449246	A	27/12/2002
				CA	2480452	A	30/10/2003
				CN	1646531	A	27/07/2005
				DE	60203619	D,T	15/09/2005
				EP	1397038	A,B	17/03/2004
				SE	1397038	T3	
				EP	1499613	A	26/01/2005
				JP	2005526836	T	08/09/2005
				NO	20044998	A	18/01/2005
				SE	0201193	D	00/00/0000
				US	20040182325	A	23/09/2004
				US	20050234036	A	20/10/2005
				AU	2002357543	A	00/00/0000
				BR	0214388	A	03/11/2004
				EP	1458541	A	22/09/2004
				JP	2005511363	T	28/04/2005
				SE	0202239	D	00/00/0000
				US	20040239005	A	02/12/2004
<hr/>							
WO	9940091	A1	12/08/1999	AU	747920	B	30/05/2002
				AU	2659099	A	23/08/1999
				CA	2319275	A	12/08/1999
				EP	1054887	A	29/11/2000
				JP	2003502272	T	21/01/2003
				US	6187777	B	13/02/2001
				US	6583154	B	24/06/2003
				ZA	9900967	A	06/08/1999
				US	6566503	B	20/05/2003
				US	20030027990	A	06/02/2003
<hr/>							
WO	0059449	A2	12/10/2000	AU	4053800	A	23/10/2000
				BR	0011182	A	10/06/2003
				CA	2367143	A	12/10/2000
				CN	1377353	A,T	30/10/2002
				EP	1169321	A	09/01/2002
				HU	0200938	A	28/10/2002
				IL	145701	D	00/00/0000
				JP	2002541078	T	03/12/2002
				US	6294541	B	25/09/2001
				US	6319928	B	20/11/2001
				US	6413975	B	02/07/2002
				US	20030073834	A	17/04/2003
<hr/>							
WO	2005037835	A1	28/04/2005	SE	0302756	D	00/00/0000
<hr/>							
WO	03000694	A1	03/01/2003	EP	1409489	A	21/04/2004
				ES	1049274	U,Y	16/11/2001
				ES	2193839	A,B	01/11/2003
				JP	2004534828	T	18/11/2004
				US	20050070558	A	31/03/2005

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/SE 2005/001835

WO	03082873	A1	09/10/2003	AU	2003215680	A	00/00/0000
				EP	1492793	A	05/01/2005
				ES	1051565	U,Y	16/08/2002
				ES	2208063	A,B	01/06/2004
				JP	2005526817	T	08/09/2005
				US	20050261248	A	24/11/2005

WO	2005042534	A2	12/05/2005	US	20050119287	A	02/06/2005
----	------------	----	------------	----	-------------	---	------------

WO	9618400	A1	20/06/1996	AT	247655	T	15/09/2003
				AU	4527996	A	03/07/1996
				CA	2206804	A,C	20/06/1996
				DE	69531555	D,T	17/06/2004
				EP	0799040	A,B	08/10/1997
				IL	116382	D	00/00/0000
				JP	2001523213	T	20/11/2001
				US	5977119	A	02/11/1999
				US	6025361	A	15/02/2000
				US	6268373	B	31/07/2001
